

APPENDIX 1

Multivariate HR estimates and 95% confidence intervals for CD by tertile of energy-adjusted protein intake, further adjusted for daily alcohol intake, physical activity, body mass index, tobacco smoking, menopause hormone therapy, or socio-economic status in the E3N cohort ($n=67,581$), 1993–2005.

	Tertile 1	Tertile 2	Tertile 3
Multivariate model (model 1)	1 (—)	4.51 (1.28–15.83)	2.70 (0.69–10.52)
Model 1+alcohol intake	1 (—)	4.47 (1.27–15.71)	2.72 (0.70–10.63)
Model 1+body mass index	1 (—)	4.52 (1.28–15.88)	2.72 (0.70–10.70)
Model 1+menopausal hormone therapy	1 (—)	4.47 (1.27–15.71)	2.68 (0.69–10.44)
Model 1+physical activity	1 (—)	4.53 (1.29–15.90)	2.72 (0.69–10.47)
Model 1+tobacco consumption	1 (—)	4.52 (1.29–15.89)	2.72 (0.69–10.50)
Model 1+socioeconomic status	1 (—)	4.51 (1.28–15.83)	2.70 (0.69–10.54)

APPENDIX 2

Multivariate HR estimates and 95% confidence intervals for UC by tertile of energy-adjusted protein intake, further adjusted for daily alcohol intake, physical activity, body mass index, tobacco smoking, menopause hormone therapy, or socio-economic status in the E3N cohort ($n=67,581$), 1993–2005.

	Tertile 1	Tertile 2	Tertile 3
Multivariate model (model 1)	1 (—)	1.62 (0.65–4.02)	3.29 (1.34–8.04)
Model 1+alcohol intake	1 (—)	1.56 (0.63–3.89)	3.06(1.25–7.50)
Model 1+body mass index	1 (—)	1.63 (0.65–4.06)	3.36 (1.36–8.29)
Model 1+menopausal hormone therapy	1 (—)	1.61 (0.65–4.00)	3.27 (1.34–8.00)
Model 1+physical activity	1 (—)	1.61 (0.65–4.00)	3.29 (1.35–8.07)
Model 1+tobacco consumption	1 (—)	1.62 (0.65–4.03)	3.26 (1.33–7.98)
Model 1+socioeconomic status	1 (—)	1.62 (0.65–4.04)	3.31(1.35–8.10)