

Appendix. References for Table 1

References:

1. Acromite MT, Mantzoros CS, Leach RE, Hurwitz J, Dorey LG. Androgens in preeclampsia. *Am J Obstet Gynecol* 1999;180:60.
2. Álvarez SI, Castañón SG, Ruata MLC, et al. Updating of normal levels of copper, zinc and selenium in serum of pregnant women. *J Trace Elem Med Biol* 2007;21:49.
3. Ardawi MSM, Nasrat HAN, BA'Aqueel HS. Calcium-regulating hormones and parathyroid hormone-related peptide in normal human pregnancy and postpartum: a longitudinal study. *Eur J Endocrinol* 1997;137:402.
4. Karim SA, Khurshid M, Rizvi JH, Jafarey SN, Rizwana I. Platelets and leucocyte counts in pregnancy. *J Pak Med Assoc* 1992;42:86–7.
5. Bacq Y, Zarka O, Bréchot JF, Mariotte N, Vol S, Tichet J, Weill J. Liver function tests in normal pregnancy: a prospective study of 102 pregnant women and 102 matched controls. *Hepatology* 1996;23:1030–4.
6. Balloch AJ, Cauchi MN. Reference ranges for haematology parameters in pregnancy derived from patient populations. *Clin Lab Haemat* 1993;15:7.
7. Beguin Y, Lipscei G, Thoumsin H, Fillet G. Blunted erythropoietin production and decreased erythropoiesis in early pregnancy. *Blood* 1991;78:89–93.

Abbassi-Ghanavati M, Greer L, Cunningham FG. Establishing normal laboratory value ranges for pregnant women. *Obstet Gynecol* 2009;114.

8. Belo L, Caslake M, Gaffney D, Santos-Silva A, Pereira-Leite L, Quintanilha A, et al. Changes in LDL size and HDL concentration in normal and preeclamptic pregnancies. *Atherosclerosis* 2002;162:425–32.
9. Belo L, Santos-Silva A, Rocha S, Caslake M, Cooney J, Pereira-Leite L, et al. Fluctuations in C-reactive protein concentration and neutrophil activation during Norman human pregnancy. *Eur J Obstet Gynecol Reprod Biol* 2005;123:46–51.
10. Bianco I, Mastropietro F, D'Asero C, Graziani B, Piergrossi P, Mezzabotta M, et al. Serum levels of erythropoietin and soluble transferrin receptor during pregnancy in non- β -thalassemic and β -thalassemic women. *Haematologica* 2000;85:902–7.
11. Borghi C, Esposti DD, Immordino V, Cassani A, Boschi S, Bovicelli L, et al. Relationship of systemic hemodynamics, left ventricular structure and function, and plasma natriuretic peptide concentrations during pregnancy complicated by preeclampsia. *Am J Obstet Gynecol* 2000;183:140–7.
12. Hauser SL, Longo DL. Harrison's Principles of Internal Medicine 15th ed . In: Braunwald E, Fauci AS, Kasper DL, Jameson JL, editors. New York (NY): McGraw-Hill; 2001. Appendices, p. A-1.
13. Carranza-Lira S, Hernández F, Sánchez M, Murrieta S, Hernández A, Sandoval C. Prolactin secretion in molar and normal pregnancy. *Int J Gynaecol Obstet* 1998;60:137–41.
14. Carter J: Serum bile acids in normal pregnancy. *Br J Obstet Gynaecol* 1991;98:540–3.

Abbassi-Ghanavati M, Greer L, Cunningham FG. Establishing normal laboratory value ranges for pregnant women. *Obstet Gynecol* 2009;114.

15. Cerneca F, Ricci G, Simeone R, Malisano M, Alberico S, Guaschino S. Coagulation and fibrinolysis changes in normal pregnancy increased levels of procoagulants and reduced levels of inhibitors during pregnancy induce a hypercoagulable state, combined with a reactive fibrinolysis. *Eur Eur J Obstet Gynecol Reprod Biol* 1997;73:31–6.
16. Choi JW, Pai SH. Tissue plasminogen activator levels change with plasma fibrinogen concentrations during pregnancy. *Ann Hematol* 2002;81:611–5. Epub 2002 Oct 23.
17. Davison JB, Vallotton MB, Lindheimer MD. Plasma osmolality and urinary concentration and dilution during and after pregnancy: evidence that lateral recumbency inhibits maximal urinary concentrating ability. *Br J Obstet Gynaecol* 1981;88:472–9.
18. Desoye G, Schweditsch MO, Pfeiffer KP, Zechner R, Kostner GM. Correlation of hormones with lipid and lipoprotein levels during normal pregnancy and postpartum. *J Clin Endocrinol Metab* 1987;64:704–12.
19. Dunlop W: Serial changes in renal haemodynamics during normal human pregnancy. *Br J Obstet Gynaecol* 1981;88:1–9.
20. Dux S, Yaron A, Carmel A, Rosenfeld JB. Renin, aldosterone, and serum-converting enzyme activity during normal and hypertensive pregnancy. *Gynecol Obstet Invest* 1984;17:252–7.
21. Elsheikh A, Creatsas G, Mastorakos G, Milingos S, Loutradis D, Michalas S. The renin-aldosterone system during normal and hypertensive pregnancy. *Arch Gynecol Obstet* 2001;264:182–5.

Abbassi-Ghanavati M, Greer L, Cunningham FG. Establishing normal laboratory value ranges for pregnant women. *Obstet Gynecol* 2009;114.

22. Ezimokhai M, Davison JM, Philips PR, Dunlop W. Non-postural serial changes in renal function during the third trimester of normal human pregnancy. *Br J Obstet Gynaecol* 1981;88:465–71.
23. Fadel HE, Northrop G, Misenheimer HR, Harp RJ. Acid-base determinations in amniotic fluid and blood of normal late pregnancy. *Obstet Gynecol* 1979;53:99–104.
24. Faught W, Garner P, Jones G, Ivey B. Changes in protein C and protein S levels in normal pregnancy. *Am J Obstet Gynecol* 1995;172:147–50.
25. Francalanci I, Comeglio P, Liotta AA, Cellai AP, Fedi S, Parretti E. D-dimer concentrations during normal pregnancy, as measured by ELISA. *Thromb Res* 1995;78:399–405.
26. Handwerker SM, Altura BT, Altura BM. Serum ionized magnesium and other electrolytes in the antenatal period of human pregnancy. *J Am Coll Nutr* 1996;15:36–43.
27. Higby K, Suiter CR, Phelps JY, Siler-Khodr T, Langer O. Normal values of urinary albumin and total protein excretion during pregnancy. *Am J Obstet Gynecol* 1994;171:984–9.
28. Hwang HS, Kwon JY, Kim MA, Park YW, Kim YH. Maternal serum highly sensitive C-reactive protein in normal pregnancy and pre-eclampsia. *Int J Gynaecol Obstet* 2007;98:105–9. Epub 2007 Jun 22.
29. Hytten FE, Lind T: Diagnostic Indices in Pregnancy. Summit, New Jersey, CIBA-GEIGY Corporation, 1975.

Abbassi-Ghanavati M, Greer L, Cunningham FG. Establishing normal laboratory value ranges for pregnant women. *Obstet Gynecol* 2009;114.

30. Ilhan N, Ilhan N, Simsek M. The changes of trace elements, malondialdehyde levels and superoxide dismutase activities in pregnancy with or without preeclampsia. *Clin Biochem* 2002;35:393–7.
31. Jimenez DM, Pocovi M, Ramon-Cajal J, Romero MA, Martinez H, Grande F. Longitudinal study of plasma lipids and lipoprotein cholesterol in normal pregnancy and puerperium. *Gynecol Obstet Invest* 1988;25:158–64.
32. Karsenti D, Bacq Y, Bréchot JF, Mariotte N, Vol S, Tichet J. Serum amylase and lipase activities in normal pregnancy: a prospective case-control study. *Am J Gastroenterol* 2001;96:697–9.
33. Kato T, Seki K, Matsui H, Sekiya S. Monomeric calcitonin in pregnant women and in cord blood. *Obstet Gynecol* 1998;92:241–4.
34. Kim EH, Lim JH, Kim YH, Park YW. The relationship between aldosterone to renin ratio and RI value of the uterine artery in the preeclamptic patient vs. normal pregnancy. *Yonsei Med J* 2008;49:138–43.
35. Kline JA, Williams GW, Hernandez-Nino J. D-dimer concentrations in normal pregnancy: new diagnostic thresholds are needed. *Clin Chem* 2005;51:825–9. Epub 2005 Mar 11.
36. Koscica KL, Bebbington M, Bernstein PS. Are maternal serum troponin I levels affected by vaginal or cesarean delivery? *Am J Perinatol* 2004;21:31–4.

Abbassi-Ghanavati M, Greer L, Cunningham FG. Establishing normal laboratory value ranges for pregnant women. *Obstet Gynecol* 2009;114.

37. Kratz A, Pesce MA, Fink DJ.: Appendix: Laboratory values of clinical importance. In: Fauci AS, Braunwald E, Kasper DL, editors. Harrison's Principles of Internal Medicine. 17th ed. New York (NY): McGraw-Hill; 2008, Appendix 1, p. A-1.
38. Larrea F, Méndez I, Parra A, Espinosa de los Monteros A. Serum pattern of different molecular forms of prolactin during normal human pregnancy. *Hum Reprod* 1993;8:1617–22.
39. Larsson A, Palm M, Hansson LO, Axelsson O. Reference values for clinical chemistry tests during normal pregnancy. *BJOG* 2008;115:874–81.
40. Lefkowitz JB, Clarke SH, Barbour LA: Comparison of protein S functional and antigenic assays in normal pregnancy. *Am J Obstet Gynecol* 1996;175:657–60.
41. Leiserowitz GS, Evans AT, Samuels SJ, Omand K, Kost GJ. Creatine kinase and its MB isoenzyme in the third trimester and the peripartum period. *J Reprod Med* 1992;37:910–6.
42. Lockitch G. Handbook of diagnostic biochemistry and hematology in normal pregnancy. Boca Raton (FL): CRC Press; 1993.
43. López-Quesada E, Vilaseca MA, Lailla JM. Plasma total homocysteine in uncomplicated pregnancy and in preeclampsia. *Eur J Obstet Gynecol Reprod Biol* 2003;108:45–9.
44. Louro MO, Cocho JA, Tutor JC. Assessment of copper status in pregnancy by means of determining the specific oxidase activity of ceruloplasmin. *Clin Chim Acta* 2001;312:123–7.

Abbassi-Ghanavati M, Greer L, Cunningham FG. Establishing normal laboratory value ranges for pregnant women. *Obstet Gynecol* 2009;114.

45. Milman N, Bergholt T, Byg KE, Eriksen L, Hvas AM. Reference intervals for haematological variables during normal pregnancy and postpartum in 434 healthy Danish women. *Eur J Haematol* 2007;79:39–46.
46. Milman N, Byg KE, Hvas AM, Bergholt T, Eriksen L. Erythrocyte folate, plasma folate and plasma homocysteine during normal pregnancy and postpartum: a longitudinal study comprising 404 Danish women. *Eur J Haematol* 2006;76:200–5. Epub 2006 Jan 12.
47. Milman N, Graudal N, Nielsen OJ, Agger AO. Serum erythropoietin during normal pregnancy: relationship to hemoglobin and iron status markers and impact of iron supplementation in a longitudinal, placebo-controlled study on 118 women. *Int J Hematol* 1997;66:159–68.
48. Mimouni F, Tsang RC, Hertzberg VS, Neumann V, Ellis K. Parathyroid hormone and calcitrol changes in normal and insulin-dependent diabetic pregnancies. *Obstet Gynecol* 1989;74:49–54.
49. Montelongo A, Lasunción MA, Pallardo LF, Herrera E. Longitudinal study of plasma lipoproteins and hormones during pregnancy in normal and diabetic women. *Diabetes* 1992;41:1651–9.
50. Moran P, Baylis PH, Lindheimer MD, Davison JM. Glomerular ultrafiltration in normal and preeclamptic pregnancy. *J Am Soc Nephrol* 2003;14:648–52.
51. Morse M. Establishing a normal range for d-dimer levels through pregnancy to aid in the diagnosis of pulmonary embolism and deep vein thrombosis. *J Thromb Haemost* 2004;2:1202–4.

Abbassi-Ghanavati M, Greer L, Cunningham FG. Establishing normal laboratory value ranges for pregnant women. *Obstet Gynecol* 2009;114.

52. O'Leary P, Boyne P, Flett P, Beilby J, James I. Longitudinal assessment of changes in reproductive hormones during normal pregnancy. *Clin Chem* 1991;37:667–72.
53. Ozerol E, Ozerol I, Gökdeniz R, Temel I, Akyol O. Effect of smoking on serum concentrations of total homocysteine, folate, vitamin B₁₂, and nitric oxide in pregnancy: a preliminary study. *Fetal Diagn Ther* 2004;19:145–8.
54. Parente JV, Franco JG Jr, Greene LJ, Meirelles RS, Martinez AR. Angiotensin-converting enzyme: serum levels during normal pregnancy. *Am J Obstet Gynecol* 1979;135:586–9.
55. Piechota W, Staszewski A. Reference ranges of lipids and apolipoproteins in pregnancy. *Eur J Obstet Gynecol Reprod Biol* 1992;45:27–35.
56. Pitkin RM, Gebhardt MP. Serum calcium concentrations in human pregnancy. *Am J Obstet Gynecol* 1977;127:775–8.
57. Price A, Obel O, Cresswell J, Catch I, Rutter S, Barik S, Heller SR, Weetman AP. Comparison of thyroid function in pregnant and non-pregnant Asian and western Caucasian women. *Clin Chim Acta* 2001;308:91–8.
58. Qvist I, Abdulla M, Jägerstad M, Svensson S. Iron, zinc and folate status during pregnancy and two months after delivery. *Acta Obstet Gynecol Scand* 1986;65:15–22.
59. Radder JK, Van Roosmalen J. HbA1c in healthy, pregnant women. *Neth J Med* 2005;63:256–9.

Abbassi-Ghanavati M, Greer L, Cunningham FG. Establishing normal laboratory value ranges for pregnant women. *Obstet Gynecol* 2009;114.

60. Reiter EO, Braunstein GD, Vargas A, Root AW. Changes in 25-hydroxyvitamin D and 24,25-dihydroxyvitamin D during pregnancy. *Am J Obstet Gynecol* 1979;135:227–9.
61. Risberg A, Larsson A, Olsson K, Lyrenäs S, Sjöquist M. Relationship between urinary albumin and albumin/creatinine ratio during normal pregnancy and pre-eclampsia. *Scand J Clin Lab Invest* 2004;64:17–23.
62. Romslo I, Haram K, Sagen N, Augensen K. Iron requirement in normal pregnancy as assessed by serum ferritin, serum transferring saturation and erythrocyte protoporphyrin determinations. *Br J Obstet Gynaecol* 1983;90:101–7.
63. Shakhmatova EI, Osipova NA, Natochin YV. Changes in osmolality and blood serum ion concentrations in pregnancy. *Hum Physiol* 2000;26:92.
64. Sharma SC, Sabra A, Molloy A, Bonnar J. Comparison of blood levels of histamine and total ascorbic acid in pre-eclampsia with normal pregnancy. *Hum Nutr Clin Nutr* 1984;38:3–9.
65. Shivvers SA, Wians FH Jr, Keffer JH, Ramin SM. Maternal cardiac troponin I levels during labor and delivery. *Am J Obstet Gynecol* 1999;180:122.
66. Singh HJ, Mohammad NH, Nila A. Serum calcium and parathormone during normal pregnancy in Malay women. *J Matern Fetal Med* 1999;8:95–100.
67. Spiropoulos K, Prodromaki E, Tsapanos V. Effect of body position on PaO₂ and PaCO₂ during pregnancy. *Gynecol Obstet Invest* 2004;58:22–5. Epub 2004 Mar 15.

Abbassi-Ghanavati M, Greer L, Cunningham FG. Establishing normal laboratory value ranges for pregnant women. *Obstet Gynecol* 2009;114.

68. Strickland DM, Hauth JC, Widish J, Strickland K, Perez R. Amylase and isoamylase activities in serum of pregnant women. *Obstet Gynecol* 1984;63:389–91.
69. Suri D, Moran J, Hibbard JU, Kasza K, Weiss RE. Assessment of adrenal reserve in pregnancy: defining the normal response to the adrenocorticotropin stimulation test. *J Clin Endocrinol Metab* 2006;91:3866–72. Epub 2006 Aug 8.
70. van Buul EJ, Steegers EA, Jongsma HW, Eskes TK, Thomas CM, Hein PR. Haematological and biochemical profile of uncomplicated pregnancy in nulliparous women; a longitudinal study. *Neth J Med* 1995;46:73–85.
71. van den Broek NR, Letsky EA. Pregnancy and the erythrocyte sedimentation rate. *BJOG* 2001;108:1164–7.
72. Walker MC, Smith GN, Perkins SL, Keely EJ, Garner PR. Changes in homocysteine levels during normal pregnancy. *Am J Obstet Gynecol* 1999;180:660–4.
73. Wickström K, Edelstam G, Löwbeer CH, Hansson LO, Siegbahn A. Reference intervals for plasma levels of fibronectin, von Willebrand factor, free protein S and antithrombin during third-trimester pregnancy. *Scand J Clin Lab Invest* 2004;64:31–40.

Abbassi-Ghanavati M, Greer L, Cunningham FG. Establishing normal laboratory value ranges for pregnant women. *Obstet Gynecol* 2009;114.