

**Table 5:** Patients Included in the Outcome Follow-Up (Greater Than or Equal to 18 Months) Analyses Compared to All Excluded Due to Inadequate (Less Than 18 Months) Follow-Up

	Benigns+EIN+Cancers Included in Analysis (n=142)	Benigns+EIN Excluded (n=35)	Fisher's Exact <i>P</i> -value
Age			
<49	43 (30.3)	15 (42.9)	0.09
49-55	51 (35.9)	6 (17.1)	
>55	48 (33.8)	14 (40.0)	
Mean (standard error)	52.8 (9.8)	53.8 (16.5)	0.71
Ethnicity			
White	70 (81.4)	11 (64.7)	0.19
Non-white	16 (18.6)	6 (35.3)	
Marital status			
Single	46 (33.3)	11 (40.7)	0.51
Married	92 (66.7)	16 (59.3)	
Insurance status			
Free care/Medicaid	20 (14.5)	8 (27.6)	0.10
Private	118 (85.5)	21 (72.4)	
Indication for initial endometrial sampling			
Postmenopausal bleeding	59 (43.1)	18 (58.1)	0.15
Heavy bleeding	29 (21.2)	4 (12.9)	
Irregular bleeding	27 (19.7)	2 (6.4)	
Other	22 (16.1)	7 (22.6)	

EIN, endometrial intraepithelial neoplasia.

Semere LG, Ko E, Johnson NR, Vitonis AF, Phang LJ et al. Endometrial intraepithelial neoplasia clinical correlates and outcomes. *Obstet Gynecol* 2011;118.

The authors provided this information as a supplement to their article.

© Copyright 2011 American College of Obstetricians and Gynecologists.