

Appendix. Births With Selected Major Congenital Anomalies Modified From the International Clearinghouse For Birth Defects Surveillance and Research Classification System Among Pregnancies With and Without Abruption From 1987–2005.

Selected Major Congenital Anomalies	Abruptio n=4,190		No Abruptio n=12,570		Adjusted*	
	n	per 10,000	n	per 10,000	OR	(95 % CI)
Anencephaly	2	5	1	1	3.57	(0.29–44.4)
Spina bifida	1	2	3	2	0.85	(0.08–9.17)
Encephalocele	1	2	0	0		
Microcephaly	1	2	4	3	0.62	(0.07–5.71)
Holoprosencephaly	0	0	2	2		
Hydrocephaly	10	24	4	3	7.16	(2.23–23.0)
Transposition of great vessels	3	7	5	4	1.91	(0.45–8.05)
Tetralogy of Fallot	5	12	5	4	3.06	(0.88–10.7)
Hypoplastic left heart syndrome	2	5	4	3	1.34	(0.24–7.59)
Coarctation of the aorta	5	12	11	9	1.33	(0.46–3.88)
Anophthalmos/microphthalmos	1	2	0	0		
Anotia/microtia	2	5	5	4	1.15	(0.22–6.18)
Choanal atresia, bilateral	0	0	2	2		
Cleft palate without cleft lip	7	17	18	14	1.08	(0.45–2.60)
Cleft lip with or without cleft palate	11	26	18	14	1.69	(0.79–3.62)
Oesophageal atresia/stenosis	14	33	9	7	4.92	(2.12–11.4)
Duodenal atresia/stenosis	4	10	2	2	6.84	(1.25–37.5)
Small intestine atresia/stenosis	1	2	1	1	3.17	(0.20–51.3)

Riihimäki O, Metsäranta M, Ritvanen A, Gissler M, Luukkaala T, Paavonen J et al. Increased prevalence of major congenital anomalies in births with placental abruption. *Obstet Gynecol* 2013;122.

The authors provided this information as a supplement to their article.

Anorectal atresia/stenosis	7	17	6	5	3.06	(1.01–9.23)
Diaphragmatic hernia	5	12	1	1	16.03	(1.86–138.4)
Omphalocele	3	7	4	3	2.12	(0.46–9.64)
Gastroschisis	1	2	2	2	1.45	(0.13–16.4)
Prune belly sequence	1	2	0	0		
Renal agenesis, bilateral	2	5	1	1	6.08	(0.54–67.8)
Cystic kidneys	6	14	6	5	2.79	(0.88–8.79)
Bladder exstrophy	2	5	1	1	5.37	(0.47–61.1)
Hypospadias, severe	11	26	5	4	7.14	(2.47–20.7)
Indeterminate sex	1	2	1	1	1.81	(0.09–34.8)
Polydactyly, preaxial	2	5	4	3	1.77	(0.32–9.70)
Limb reduction defects	5	12	8	6	1.95	(0.63–6.00)
Trisomy 13	1	2	0	0		
Trisomy 18	3	7	7	6	1.27	(0.32–5.10)
Down Syndrome	3	7	15	12	0.58	(0.17–2.03)

*Adjusted by maternal age, parity, socioeconomic status and smoking during pregnancy.
A limit for statistical significance in this table was $P<.01$.

Riihimäki O, Metsäranta M, Ritvanen A, Gissler M, Luukkaala T, Paavonen J et al. Increased prevalence of major congenital anomalies in births with placental abruption. *Obstet Gynecol* 2013;122.

The authors provided this information as a supplement to their article.