

**Appendix 3. Distribution of Stroke Type by Hypertensive Disorders of Pregnancy Status and Pregnancy Period: Pregnancy Hospitalizations, Nationwide Inpatient Sample, 1994 to 2011**

	<b>Overall (N = 31,616) n (%)</b>			<b>Hospitalizations With HDP- Associated Stroke (n = 9,877) n (%)</b>			<b>Hospitalizations With Non-HDP- Associated Stroke (n = 21,740) n (%)</b>		
<b>By Stroke Subtype</b>	<i>Antenatal</i>	<i>Delivery</i>	<i>Postpartum</i>	<i>Antenatal</i>	<i>Delivery</i>	<i>Postpartum</i>	<i>Antenatal</i>	<i>Delivery</i>	<i>Postpartum</i>
<i>Hemorrhagic Stroke</i>	2,428 (24.0)	2,093 (19.4)	4,255 (39.4)	307 (21.2)	1,188 (28.0)	1,911 (45.6)	2,121 (24.5)	905 (13.9)	2,344 (35.5)
<i>Ischemic Stroke*</i>	5,554 (55.0)	3,062 (28.5)	3,309 (30.6)	839 (57.8)	1,394 (32.8)	1,178 (28.1)	4,715 (54.5)	1,668 (25.6)	2,131 (32.2)
<i>Unspecified</i>	2,106 (21.0)	5,594 (52.0)	3,217 (30.0)	300 (21.0)	1,667 (39.2)	1,093 (26.3)	1,806 (21.0)	3,927 (60.5)	2,123 (32.3)
<i>Combined Stroke</i>	10,086	10,748	10,781	1,446	4,249	4,182 (42.3)	8,642	6,500	6,598 (30.4)

Leffert LR, Clancy CR, Bateman BT, Bryant AS, Kuklina EV. Hypertensive disorders and pregnancy-related stroke: frequency, trends, risk factors, and outcomes. *Obstet Gynecol* 2014;125.

The authors provided this information as a supplement to their article.

© Copyright 2014 American College of Obstetricians and Gynecologists.

<i>Subtypes</i>	(31.9)	(34.0)	(34.1)	(14.7)	(43.0)		(39.7)	(29.9)	
<p>* Ischemic stroke group includes ischemic stroke and transient ischemic attack</p> <p>The total percentage may be not summing to 100% due to rounding.</p>									