

## Appendix 1. Acknowledgements

Institutions involved in the Consortium include, in alphabetical order: Baystate Medical Center, Springfield, MA; Cedars-Sinai Medical Center Burnes Allen Research Center, Los Angeles, CA; Christiana Care Health System, Newark, DE; Georgetown University Hospital, MedStar Health, Washington, DC; Indiana University Clarian Health, Indianapolis, IN; Intermountain Healthcare and the University of Utah, Salt Lake City, Utah; Maimonides Medical Center, Brooklyn, NY; MetroHealth Medical Center, Cleveland, OH.; Summa Health System, Akron City Hospital, Akron, OH; The EMMES Corporation, Rockville MD (Data Coordinating Center); University of Illinois at Chicago, Chicago, IL; University of Miami, Miami, FL; and University of Texas Health Science Center at Houston, Houston, Texas. The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the *Eunice Kennedy Shriver* National Institute of Child Health and Human Development, National Institutes of Health.

Halscott TL, Reddy UM, Landy HJ, Ramsey PS, Iqbal SN, Huang CC, et al. Maternal and neonatal outcomes by attempted mode of operative delivery from a low station in the second stage of labor. *Obstet Gynecol* 2015;126.

The authors provided this information as a supplement to their article.