

Appendix.

In addition to the authors, the following Pelvic Floor Disorders Network Members participated in the CARE trial: *University of Alabama at Birmingham, Birmingham, Alabama* – K. Burgio, P. Goode, R. Varner, V. Willis; *Baylor College of Medicine, Houston, Texas* – P. Thompson, P. Lotze, N. Frierson; *University of Iowa, Iowa City, Iowa* – D. Brandt, D. Haury, K. Kreder, C. Bradley; *Johns Hopkins Medical Institutes, Baltimore, Maryland* – V. Handa, R. Gutman, M. Sauter; *Loyola University, Chicago, Illinois* – M. FitzGerald, K. Kenton, D. Koch, C. Ball; *University of Michigan, Ann Arbor, Michigan* – J. Wei, J. DeLancey, N. Janz, D. Smith, P. Wren, J. Imus, B. Marchant, Y. Casher; *University of North Carolina at Chapel Hill, Chapel Hill, North Carolina* – A. Connolly, M. Jannelli, M. Loomis, A. Murphy, E. Wells, W. Whitehead; *University of Pittsburgh/Magee-Women's Hospitals, Pittsburgh, Pennsylvania* – D. Borello-France, J. Gruss, W. Leng, P. Moalli, H. Zyczynski

Online appendix to Brubaker L, Nygaard I, Richter HE, Visco A, Weber AM, Cundiff GW et al. Two-year outcomes after sacrocolpopexy with and without burch to prevent stress urinary incontinence. *Obstet Gynecol* 2008;112:49–55.

© 2008 by The American College of Obstetricians and Gynecologists.