

Appendix 1. Severe Maternal Morbidity Indicators by Percent of Deliveries

Severe Maternal Morbidity Indicator	N	Percent of Deliveries	Percent of Severe Maternal Morbidity*
1. Acute myocardial infarction	12	0.003%	0.1%
2. Acute renal failure	242	0.10%	2.7%
3. Adult respiratory distress syndrome	272	0.08%	3.1%
4. Amniotic fluid embolism	15	0.004%	0.2%
5. Aneurysm#	--	--	--
6. Cardiac arrest/ventricular fibrillation	24	0.01%	0.3%
7. Disseminated intravascular coagulation	1605	0.45%	18.1%
8. Eclampsia	183	0.05%	2.1%
9. Heart failure during procedure or surgery	1102	0.31%	12.4%
10. Internal injuries of thorax, abdomen, and pelvis	11	0.003%	0.1%
11. Intracranial injuries#	--	--	--
12. Puerperal cerebrovascular disorders	152	0.04%	1.7%
13. Pulmonary edema	117	0.03%	1.3%
14. Severe anesthesia complications	104	0.03%	1.2%
15. Sepsis	124	0.04%	1.4%
16. Shock	187	0.05%	2.1%
17. Sickle cell anemia with crisis	80	0.02%	0.9%

Howell EA, Egorova NN, Janevic T, Balbierz A, Zeitlin J, Hebert PL. Severe maternal morbidity among hispanic women in New York City: investigation of health disparities. *Obstet Gynecol* 2017; 129.

The authors provided this information as a supplement to their article.

18. Thrombotic embolism	103	0.03%	1.2%
19. Blood transfusion	7120	2.01%	80.2%
19a. Blood transfusion only	6037	1.71%	67%
20. Cardio monitoring	61	0.02%	0.7%
21. Conversion of cardiac rhythm	38	0.01%	0.4%
22. Hysterectomy	504	0.14%	5.7%
23. Operations on heart and pericardium	259	0.07%	2.9%
24. Temporary tracheostomy#	--	--	--
25. Ventilation	425	0.12%	4.8%
26. Deaths	31	0.01%	0.3%

Howell EA, Egorova NN, Janevic T, Balbierz A, Zeitlin J, Hebert PL. Severe maternal morbidity among hispanic women in New York City: investigation of health disparities. *Obstet Gynecol* 2017; 129.

The authors provided this information as a supplement to their article.