

**Appendix 2. Oxycodone Utilization After Discharge and Satisfaction With Pain Regimen, Stratified by Type and Timing Of Cesarean Delivery**

Outcomes	Median [IQR], or n (%)			
	Primary cesarean delivery (n=35)	Repeat cesarean delivery (n=15)	Planned <sup>a</sup> cesarean delivery (n=19)	Unplanned cesarean delivery (n=31)
<b>Number of oxycodone tablets chosen,</b>	20.0 [15.0, 25.0] <sup>b</sup>	20.0 [10.0, 30.0] <sup>c</sup>	20.0 [10.0, 23.0] <sup>d</sup>	20.0 [20.0, 30.0] <sup>e</sup>
<b>Number of oxycodone tablets used</b>	16.0 [8.0, 25.0]	15.0 [3.0, 30.0]	11.0 [3.0, 23.0]	16.0 [8.0, 31.0]
<b>Number of oxycodone tablets remaining</b>	4.0 [0.0, 11.0]	0.0 [0.0, 8.0]	1.0 [0.0, 9.0]	4.0 [0.0, 8.0]
<b>Need for oxycodone refills</b>	2 (5.7)	2 (13.3)	0 (0.0)	4 (12.9)
<b>Satisfied with outpatient pain management</b>	18 (51.4)	8 (53.3)	9 (47.4)	17 (54.8)
<b>Very satisfied with outpatient pain management</b>	13 (37.1)	6 (40.0)	8 (42.1)	11 (35.5)

<sup>a</sup> Planned cesarean delivery refers to a scheduled cesarean delivery. Unplanned cesarean delivery refers to an intrapartum cesarean delivery.

<sup>b</sup> For 5 patients, the number of tablets prescribed was different from the number chosen, such that the overall median [IQR] number of tablets dispensed was 23.0 [20.0, 35.0].

<sup>c</sup> For 1 patient, the number of tablets prescribed was different from the number chosen, such that the overall median [IQR] number of tablets dispensed was 20.0 [10.0, 30.0].

<sup>d</sup> For 4 patients, the number of tablets prescribed was different from the number chosen, such that the overall median [IQR] number of tablets dispensed was 20.0 [10.0, 25.0].

<sup>e</sup> For 2 patients, the number of tablets prescribed was different from the number chosen, such that the overall median [IQR] number of tablets dispensed was 25.0 [20.0, 35.0].