Appendix 1. Full Search Strategies for Electronic Databases

PubMed search:

	Search Algorithm	Search Results
#1	("leiomyoma" [Mesh] OR "leiomyomas" [tiab] OR "leiomyomata" [tiab] OR "fibroid" [tiab] OR "fibroids" [tiab] OR "myoma" [tiab] OR "myomas" [tiab] OR (fibroma [mh] AND (uterine diseases [mh] OR uterus [mh])) AND ("Pregnancy" [Mesh] OR "pregnancy" [tiab] OR "fertility" [Mesh] OR "abortion, spontaneous" [Mesh] OR "fetal death" [Mesh] OR "spontaneous abortion" [tiab] OR "miscarriage" [tiab])	3,101
#2	#1 AND ("humans"[MeSH Terms] AND English[lang])	1,971
#3	newspaper article[pt]	0
#4	letter[pt]	59
#5	comment[pt]	11
#6	case reports[pt]	514
#7	review[pt]	325
#8	practice guideline[pt]	8
#9	news[pt]	0
#10	editorial[pt]	4
#11	historical article[pt]	3
#12	meta-analysis[pt]	3
#13	legal cases[pt]	0
#14	#2 NOT (#4 OR #5 OR #6 OR #7 OR #8 OR #9 OR #10 OR #11 OR #12 OR #13)	1045

Web of Science search:

	Search Algorithm	Search Results
#1	Fibroids OR Leiomyoma OR Leiomyomata OR Myoma	23,294
#2	Miscarriage OR Spontaneous Abortion OR Abortion OR Fetal Death OR Pregnancy Loss	136,640
#3	#1 AND #2	836
#4	#3 AND English[lang]	550
#5	#4 AND article[pt]	469
#6	#5 AND "humans"	460

Sundermann AC, Velez Edwards DR, Bray MJ, Jones SH, Latham SM, Hartmann KE. Leiomyomas in pregnancy and spontaneous abortion: a systematic review and meta-analysis. Obstet Gynecol 2017; 130.

EMBASE search

	Search Algorithm	Search Results
#1	leiomyoma.mp. or exp leiomyoma/	18,886
#2	exp uterus myoma/ or leiomyomata.mp.	12,045
#3	fibroid.mp.	4,506
#4	exp myoma/ or myoma.mp.	15,703
#5	fibroma.mp. or exp fibroma/	15,133
#6	uterus.mp. or exp uterus/	197,559
#7	#5 AND #6	793
#8	#1 OR #2 OR #3 OR #4 OR #7	32,204
#9	pregnancy.mp. or exp pregnancy/	755,053
#10	exp fertility/ or fertility.mp.	102,499
#11	spontaneous abortion.mp. or exp spontaneous abortion/	33,739
#12	miscarriage.mp.	12,392
#13	fetal death.mp. or exp fetus death/	34,196
#14	pregnancy loss.mp.	6748
#15	#9 OR #10 OR #11 OR #12 OR #13 OR #14	838,294
#16	#8 AND #15	4,741
#17	limit #16 to human	4212
#18	limit #17 to english language	3375
#19	limit #18 to article	1867
#20	limit #19 to exclude MedLine journals	273
#21	#20 NOT ("case report" [pt])	170

Sundermann AC, Velez Edwards DR, Bray MJ, Jones SH, Latham SM, Hartmann KE. Leiomyomas in pregnancy and spontaneous abortion: a systematic review and meta-analysis. Obstet Gynecol 2017; 130.

The authors provided this information as a supplement to their article.

^{©2017} American College of Obstetricians and Gynecologists.

ClinicalTrials.gov

	Search Algorithm	Search Results
#1	("leiomyoma"OR "leiomyomas" OR "leiomyomata" OR "fibroid" OR	7
	"fibroids" OR "myoma" OR "myomas") AND ("pregnancy loss" OR	
	"fetal death" OR "spontaneous abortion" OR "miscarriage")	
#2	limit #1 to completed	1

Appendix 2. Data Extraction Tool

Abstract number	
First author	
Year published	
Journal	_
Study design:	Prospective CohortRetrospective CohortCase-Control
Years of study:	
Study setting:	
Describe recruitment of exposed (women with fibroids)	
Describe recruitment of unexposed (women without fibroids)	
Describe study population exclusion criteria	
How and when was fibroid status assessed in exposed group?	
How and when was fibroid status assessed in unexposed group?	
How was miscarriage defined?	
Age distribution of study population	
Distribution of race in the study population	

Sundermann AC, Velez Edwards DR, Bray MJ, Jones SH, Latham SM, Hartmann KE. Leiomyomas in pregnancy and spontaneous abortion: a systematic review and meta-analysis. Obstet Gynecol 2017; 130.

The authors provided this information as a supplement to their article.

Distribution of parity in the study population	
Distribution of induced abortion history the study population	
Distribution of BMI in the study population	
Distribution of alcohol use in the study population	
Number of women with fibroids who had a spontaneous abortion	
Number of women with fibroids who did not have a spontaneous abortion	
Number of women without fibroids who had a spontaneous abortion	
Number of women without fibroids who did not have a spontaneous abortion	
Crude point estimate and 95% confidence intervals	
Adjusted estimate provided	YesNo
Covariates included in adjusted analysis	
Adjusted point estimate and 95% confidence intervals	
Describe other analyses of the relationship between fibroids and miscarriage Risk of Bias Estimator (Cohort)	
Mak of bias Estillator (COHOLL)	

 Truly representative (one star) Somewhat representative of general population of reproductive-age women (one star) Selected group No description of derivation of the cohort
 Drawn from the same community as the exposed cohort (one star) Drawn from a different source No description of the derivation of the non-exposed cohort
 Secure record (e.g., ultrasound verified diagnosis) (one star) Diagnosis based on symptoms Self-report No description Other
o Yes o No
 The study controls for age (one star) Study controls for other factors (one star) Cohorts are not comparable on the basis of the design or analysis controlled for confounders

Sundermann AC, Velez Edwards DR, Bray MJ, Jones SH, Latham SM, Hartmann KE. Leiomyomas in pregnancy and spontaneous abortion: a systematic review and meta-analysis. Obstet Gynecol 2017; 130.

The authors provided this information as a supplement to their article.

 $\hbox{@2017 American College of Obstetricians and Gynecologists.}$

Outcome

Assessment of outcome

- Independent blind assessment (one star)
- o Record linkage (one star)
- o Self-report
- No description
- o Other

Was follow-up long enough for outcome to occur?

YesNo


Indicate the median duration of followup and a brief rationale for the assessment above

Adequacy of follow-up cohorts

- Complete follow-up: all subjects accounted for (one star)
- Subjects lost to follow-up unlikely to introduce bias: number lost less than or equal to 20% or description of those lost suggested no difference than those followed (one star)
- Follow-up rate less than 80% and no description of those loss
- No description

Appendix 3. Risk of bias assessment.

Newcastle-Ottawa Scale for risk of bias assessment of included studies¹



Abbreviations: G, good; P, poor

References:

- 1. Wells GA, Shea B, O'Connell D, Peterson J, Welch V, Losos M, et al. The Newcastle-Ottawa Scale for assessing the quality of non-randomised studies in meta-analyses. Available at: http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp. Retrieved May 15, 2016.
- 2. Viswanathan M, Ansari MT, Berkman ND, Chang S, Hartling L, McPheeters LM, et al. Assessing the

Sundermann AC, Velez Edwards DR, Bray MJ, Jones SH, Latham SM, Hartmann KE. Leiomyomas in pregnancy and spontaneous abortion: a systematic review and meta-analysis. Obstet Gynecol 2017; 130.

The authors provided this information as a supplement to their article.

^{*} Adjustment deemed adequate if study adjusted for both maternal age and maternal race

[†] Based on Agency for Healthcare Research and Quality grades: Good, Fair, Poor²

Risk of Bias of Individual Studies in Systematic Reviews of Health Care Interventions. Agency for Healthcare Research and Quality Methods Guide for Comparative Effectiveness Reviews. AHRQ Publication No 12-EHC047-EF 2012.