OBSTETRICS & GYNECOLOGY

NOTICE: This document contains correspondence generated during peer review and subsequent revisions but before transmittal to production for composition and copyediting:

- Comments from the reviewers and editors (email to author requesting revisions)
- Response from the author (cover letter submitted with revised manuscript)*

**The corresponding author has opted to make this information publicly available.*

Personal or nonessential information may be redacted at the editor's discretion.

Questions about these materials may be directed to the *Obstetrics & Gynecology* editorial office:
obgyn@greenjournal.org.

Date: Apr 12, 2019
To: "Jill Melissa Hagey" [REDACTED]
From: "The Green Journal" em@greenjournal.org
Subject: Your Submission ONG-19-377

RE: Manuscript Number ONG-19-377

Gabapentin for peri-operative pain management for uterine aspiration: a randomized controlled trial

Dear Dr. Hagey:

Your manuscript has been reviewed by the Editorial Board and by special expert referees. Although it is judged not acceptable for publication in Obstetrics & Gynecology in its present form, we would be willing to give further consideration to a revised version.

If you wish to consider revising your manuscript, you will first need to study carefully the enclosed reports submitted by the referees and editors. Each point raised requires a response, by either revising your manuscript or making a clear and convincing argument as to why no revision is needed. To facilitate our review, we prefer that the cover letter include the comments made by the reviewers and the editor followed by your response. The revised manuscript should indicate the position of all changes made. We suggest that you use the "track changes" feature in your word processing software to do so (rather than strikethrough or underline formatting).

Your paper will be maintained in active status for 21 days from the date of this letter. If we have not heard from you by May 03, 2019, we will assume you wish to withdraw the manuscript from further consideration.

REVIEWER COMMENTS:

Reviewer #1: This is an original research study describing a randomized controlled trial of adjunctive gabapentin for pain control during surgical abortion. The authors added 600mg gabapentin to their standard "minimal sedation" regimen and evaluated pain control during and at one day post procedure as well as anxiety, side effects and patient satisfaction. The authors found no difference in their primary outcome of pain at five minutes post procedure but did note significantly reduced opioid use post-procedure at one day. Importantly, the authors note widely varied practice in pain regimens post-surgical abortion and that gabapentin is a safe, well-tolerated medication that could reduce the need for opioid after a surgical abortion. Their study adds to the growing literature supporting use of gabapentin for abortion-related pain.

The study was IRB approved and registered on clinical trials.gov. The CONSORT checklist is included in the submission.

ABSTRACT:

1. Introduction: This is clear and defines up front what "usual" oral pain management means in your setting to set the context for readers.
2. Methods: In line 38, I would add that you also measured patient satisfaction with pain control at one day post procedure.
3. Results: In line 42-43, it would be important here to indicate that 73% of the participants received a narcotic prescription for post procedure pain control and that this wasn't standardized as part of the study protocol.
4. Conclusion: Line 45-46, this seems a bit vague to me here and in the discussion. I'm assuming this means studying gabapentin in local-only patients, or the effect on doses of IV opioids but if there is a way to make this more clear, that would be helpful to show the readers the implications of this study.
5. PRECIS: Need to add post-procedure opiate use since participants all still received pre-procedure opioid.

INTRODUCTION:

6. In line 52, add in some examples of what else has been studied that you are referencing here. The generalist reader may not be familiar with this literature.
7. Line 59, you start the introduction talking about surgical abortion in the US or North America but then reference uterine aspiration globally, do you mean internationally or globally like common? This is a really good point about using gabapentin for other similar gyn procedures but I was distracted by this.
8. Line 64, I would indicate you also measured patient satisfaction with the experience and pain control

MATERIALS AND METHODS:

9. Important that you included patients with chronic pain and/or history of drug alcohol use to make this result

generalizable.

10. Line 98, did you do any cervical prep for patients? If so how did that impact when/how they received their study med?
11. Line 103, how long do patients in your clinic sit with lorazepam? Was the 1-2 hours for this study different?
12. How did you decide on the five minutes after speculum removal pain score as the primary outcome? Is the standard time point the other pain studies cited use?
13. Line 124, was there any standardization of the post-operative medications for the study or was it provider discretion? Do you have any providers who prescribe only NSAIDs post-procedure?

RESULTS:

14. What percentage of your patients in this setting receive moderate IV sedation vs. oral pain management described in this study? Do you have patients that are local-only?
15. Line 165, it is very impressive only one participant didn't complete the 24 hour phone call!
16. Can you provide the breakdown of indication for abortion by group either in Table 1 or in the paragraph lines 167-170?
17. In line 193 on the influence of time, is this a result of the time they received the study med or the other oral sedation meds?
18. For women who received an opiate prescription, what was the dose and range of number given?
19. What were the results for patient satisfaction and were those different by group?

DISCUSSION:

20. Line 210 is too long for the first line of the discussion. The information in the second sentence should lead that paragraph.
21. Line 221 and line 223 seem to contradict each other. It is important to note that other studies have found that gabapentin impacted pain but what about uterine aspiration pain or the way you measured pain is different than those studies.
22. Paragraph starting with line 234 is an important finding but needs background in the methods about how prescribing is done in your setting.
23. The study in line 244 used pregabalin, this is an important distinction as pregabalin is much more expensive than gabapentin thus many sites have not adopted this evidence into practice.
24. Is there a reference for line 249 regarding the variations in clinical practice around routine prescribing of opioids post abortion?
25. A strength I would include is that you measured baseline pain, history of opiate use and sexual trauma.
26. In line 259, that data isn't presented in your results, readers can determine generalizability based on the demographics in Table 1 and how similar the protocol for oral sedation followed is to their clinical practice. While similar between study arms, having 31% of participants with a master's degree and reporting a household income of \$ >100,000 should probably be remarked on in the discussion and reflect the setting the study was done in perhaps.
27. The last paragraph should be strengthened, I would be clear in the finding that gabapentin wasn't associated with decreased pain at 5 minutes after oral sedation for first trimester aspiration abortion but was well-tolerated and led to reduced opiate use post-procedure which is what this study adds to the current literature. If you have recommendations for further research or plan to evaluate this relationship further, add that in too.

TABLES AND FIGURES

28. Table 1, can you add in indication for abortion?
29. Table 4, can you add in nausea/vomiting outcomes?

Reviewer #2:

TITLE: Gabapentin for uterine aspiration

Overall: This is a randomized controlled trial comparing the addition of gabapentin to placebo for pain management during and after uterine aspiration procedures for induced abortion. 96 women were enrolled. The study was conducted at a single institution.

Overall in the abstract, and elsewhere in the paper there is a jump from gabapentin related pain score and opioid use. This jump is implicit and not an actual measurement. Please clarify the measures in this study - VAS and gabapentin or opioid use.

Many physicians do not use opioids at all for post-op pain management after first trimester uterine evacuations so please be clear when you use it and when you don't. This paper should not ultimately serve as justification for increased use of post-operative opioids for a very minor procedure.

Does abstract provides good information but there are some places where the abstract could be stronger:

1. The objective written in the abstract should be the same as the objective and primary outcome - which is pain 5

minutes post procedure.

2. For participant flow please state the overall number screened, enrolled and included in the analysis groups per the CONSORT guidelines recommendation. Providing the screened number matches your opening statement in the results which includes the statement about the women who were excluded from enrolling.

3. Overall in the abstract, and elsewhere in the paper there is an jump from gabapentin related pain score and opioid use. This jump is implicit and not an actual measurement.

How this paper fits in with existing literature:

Author Conflicts are stated. The paper will be presented at ACOG annual meeting in May.

IRB/Ethics approval was obtained from Duke IRB

If RCT was it registered? Yes, First Posted: April 1, 2016; Last Update: November 8, 2018, as completed. Primary outcome listed as pain score on 11-point VAS 5 minutes post procedure.

The outcomes measures and intended sample on Clinical Trials site agree with those in paper.

If the CONSORT guidelines were used please state such and reference them in the paper.

CONSORT Statement: David Moher, Sally Hopewell, Kenneth F. Schulz, Victor Montori, Peter C. Gøtzsche, P.J. Devereaux, Diana Elbourne, Matthias Egger, Douglas G. Altman, CONSORT 2010 Explanation and Elaboration: updated guidelines for reporting parallel group randomised trials, Journal of Clinical Epidemiology, Volume 63, Issue 8, August 2010, Pages e1-e37, ISSN 0895-4356, <http://dx.doi.org/10.1016/j.jclinepi.2010.03.004>. (<http://www.sciencedirect.com/science/article/pii/S0895435610001034>)

TITLE: the title is good - succinct and meets CONSORT guidelines.

PRECIS: State the precis in active language.

INTRODUCTION:

1. The introduction could be a little more specific about the procedure (first trimester vacuum aspiration) and usual pain management and then the role of this with the opioid epidemic - if it fits. Overall the references to the opioid epidemic seem a bit of a tangent.
2. Lines 58: Please include in the introduction an overview or statement of how frequently opioids are actually used for postoperative pain management after first trimester uterine evacuations. Certainly in this reviewers
3. Need to be specific - uterine aspiration or dilation and evacuation.
4. Line 62: Please be clear - if you assessed please be clear which is the primary objective and which is secondary. The objective should match the outcomes and the reporting.

MATERIALS AND METHODS:

5. Overall the methods could be more concise.
6. The section on outcomes should be concise and focus on the primary and secondary outcomes. Currently there is so much information in the paragraph the details and focus is hard to follow.
7. Lines 92-94: Please state who prepared the blinding for the study medications.
8. Line 97: Headings for sections aren't needed if strong enough topic sentences are included.
9. An overall statement of the type of study should be made at the beginning of the Materials and Methods section - cross-sectional? Retrospective or prospective cohort?
10. Is the population and setting clearly stated
11. The population and setting should be clearly stated - what are the demographics of patients and clinicians at the setting the study took place?
12. The subject selection is clearly stated?
13. concise with a list of eligibility criteria only - this present the combination of positives and negatives being listed.
14. Does the process minimize bias?
15. The primary outcome is clearly stated:
16. Adverse events appropriately collected?
17. Is sample size justified? For which outcome?

RESULTS:

18. A participant flow chart should be included - this will provide readers with an understanding of the final analysis population, and this is also a crucial component of reporting according to STROBE.
19. Please include a statement of how many women were approached/screened/enrolled in the opening of the results.
20. Is participant flow chart included?
21. Response rate/follow-up rate good enough?
22. Is bias minimized? Is confounding considered?
23. How are dropouts defined and analyzed?

DISCUSSION:

24. Please state the overall findings of your study in the context of your population and setting - this study is limited in

that it's single institution.

25. The fact that MORE opiate was used by gabapentin group than placebo group actually demonstrates that there may be some harm when gabapentin is used - this must be reported and addressed in the discussion.

26. Overall the discussion could be a bit shorter.

27. Lines 234-243 - this is important and is related to comments about the overall paper - implying that gabapentin may reduce opiate use...I think this just detracts from the overall paper. What was the rationale for the study? To decrease opiate use or to decrease pain associated with uterine aspiration in first trimester for induced abortion?

28. Are limitations discussed? Are any insurmountable?

29. Is conclusion consistent with results?

TABLES AND FIGURES

30. The tables should have more descriptive figures. The tables would be easier to read esp Table 1 if the percentage signs were removed, numbers were lined up, and percentages were rounded.

31. There should be a table that clearly demonstrates opiate use during and after the procedure by group since this was a secondary outcome, and a lot of text is devoted to this concept.

Reviewer #3: Thank you for your submission.

Please see comments below:

1. Appreciate the inclusion of women with chronic pain in the study as excluding them would have decreased the ability to generalize the results.

2. Would appreciate more information regarding what the "perioperative clearance" entailed

3. The 24hr pain scale used was the reverse of the traditional pain scale used in across medical specialties (where 0=no pain and 10=most severe) which could potentially confuse patients (or even data collectors).

STATISTICAL EDITOR COMMENTS:

The Statistical Editor makes the following points that need to be addressed:

lines 112-113, 132-136: The primary outcome was specifically cited as VAS on a 100-mm scale at 5 minutes after speculum removal. The primary outcome should first be reported in the same format. That is, as two VAS 100 point VAS scales for the gabapentin vs control cohorts and cited as means(SD) with stats testing. If the distributions were more appropriately non-parametrically tested as median(Range or IQR), that could then be cited, but not as the primary outcome. Likewise, the correction for the baseline pain scale, although of interest, was not the primary outcome. Furthermore, the cohorts were randomized, so any difference in baseline pain scores would be attributable to random chance. That is, the primary outcome should be clearly demarcated from the other, secondary outcomes

General: The precis, abstract and results sections need to be re-written, so as to emphasize that the primary outcome of this RCT was negative. Some secondary outcomes were statistically significant, but those were not how the hypothesis was stated nor how the power/sample size calculated.

Table 1: Need units for age, income (annual in dollars)

Table 3: Were the anxiety scores normally distributed? If not, should cite as median(range or IQR) and test non-parametrically.

EDITOR COMMENTS:

1. Thank you for your submission to Obstetrics & Gynecology. In addition to the comments from the reviewers above, you are being sent a notated PDF that contains the Editor's specific comments. Please review and consider the comments in this file prior to submitting your revised manuscript. These comments should be included in your point-by-point response cover letter.

The notated PDF is uploaded to this submission's record in Editorial Manager. If you cannot locate the file, contact Randi Zung and she will send it by email - rzung@greenjournal.org.

My major concern for you to address in your revision is that this is essentially a negative study. For your primary outcome, there was no difference between gabapentin and placebo. Your paper as written sort of glosses over that and instead focuses on the issue of side effects and post op opioid use, which were secondary outcomes. The use of opioids post these procedures should be quite low to begin with in the average patient. The focus on the positive secondary outcomes, while not focusing on the negative primary outcome is a form of what is known as "Spin" and needs to be corrected to present this as a negative outcome, with the secondary, exploratory outcomes worth studying further.

- The Journal style doesn't not use the virgule (/) except in numeric expressions. Please edit here and in all instances.

- In both the abstract and the paper, please provide absolute numbers as well as which ever effect size you are reporting (if appropriate) + Confidence intervals. P values may be omitted for space concerns. We strongly prefer CI's as they give more information about strength of association than do P values. By absolute values, I mean something like xx (outcome in exposed)/yy(outcome in unexposed) (zz%) (Effect size= ; 95% CI=.) An example might be: Outcome 1 was more common in the exposed than the unexposed 60%/20% (Effect size=3; 95% CI 2.6-3.4)

- When you write that a study occurred between date 1 and date 2, it literally excludes those boundary dates. For instance, "This study was performed between Feb 2018 and Jan 2019" would mean it was performed from March 2018 to Dec 2018. Do you instead mean that the study was performed from date 1 to date 2? If so, please edit. The same is true on line 32 in description of the included EGA's.

- plural: procedures

- Journal style uses cesarean birth or delivery but not cesarean section.

- It is an idiosyncratic fact that at the Journal we tend to avoid the use of the word impact to imply the result of a change, preferring to limit "impact" to mean a physical blow.

- For clarity, in some places you state inclusion was of patients SCHEDULED to undergo surgical abortion and here you say it was presentation. As these are potentially different #'s of women, which is it? Were the randomized at time of scheduling or at presentation?

- These semi-colons should be commas.

- who approached them?

- is this on the day of surgery?

- Could you combine the information on lines 106-108 and 113-116?

- You were doing 3 different VAS measurements at each of these time points on a patient w/ narcotics and a anxiolytic on board. This reasonably could have taken more than 1-2 minutes to perform. Did you standardize the order of administration? Since it was. your primary end point, was the 5 minute post speculum removal pain score done first, before the anxiety and nausea scores?

- what is your standard prescription for women to go home with for pain management? Did it differ by EGA?

- For an RCT, is is unusual to have exactly equal numbers

- please state this the way you do in methods "five minutes after removal of speculum" to avoid confusion,

- are these different? Please provide statistical information as noted above in the abstract comments

- We do not allow authors to describe variables or outcomes in terms that imply a difference (such as of the terms "trend" or "tendency" or "marginally different") unless there is a statistical difference. Please edit here and throughout. Here you should start the sentence by saying "Pain five minutes after removal of the speculum, controlling for baseline pain, was not statistically different from the nul. (xxxx) with only a 3.4 mm difference. Same true for sentence on line 176

- Are these reversed? This is opposite of the design of a normal pain scale. Could that have been a source of confusion for patients?

- please provide the absolute numbers as well.

- Please be very careful about using "after the procedure" as it is a bit vague. Line 181 you talk about "on removal of the speculum and end of the procedure"--I assumed those were 2 different times. Here on line 189 you say "5 minutes after the procedure"....was that at speculum removal or some other time.

- was this the same for both groups?

- hope in discussion you comment on ability to decrease prescription for opioids overall, independent of whether you continue to use gabapentine.

- This is called a primacy claim: yours is the first, biggest, was innovative, etc...In order to assert that, you need to provide the search terms used and the data base (s) searched (PubMed, Google Scholar, etc) to substantiate the claim. Otherwise, it needs to be deleted.

- I thought you had one who did not follow up at 24 hours? (Line 165)

- excellent graphic

2. The Editors of Obstetrics & Gynecology are seeking to increase transparency around its peer-review process, in line with efforts to do so in international biomedical peer review publishing. If your article is accepted, we will be posting this revision letter as supplemental digital content to the published article online. Additionally, unless you choose to opt out, we will also be including your point-by-point response to the revision letter, as well as subsequent author queries. If you opt out of including your response, only the revision letter will be posted. Please reply to this letter with one of two responses:

- a. OPT-IN: Yes, please publish my response letter and subsequent email correspondence related to author queries.
- b. OPT-OUT: No, please do not publish my response letter and subsequent email correspondence related to author queries.

3. As of December 17, 2018, Obstetrics & Gynecology has implemented an "electronic Copyright Transfer Agreement" (eCTA) and will no longer be collecting author agreement forms. When you are ready to revise your manuscript, you will be prompted in Editorial Manager (EM) to click on "Revise Submission." Doing so will launch the resubmission process, and you will be walked through the various questions that comprise the eCTA. Each of your coauthors will receive an email from the system requesting that they review and electronically sign the eCTA.

Any author agreement forms previously submitted will be superseded by the eCTA. During the resubmission process, you are welcome to remove these PDFs from EM. However, if you prefer, we can remove them for you after submission.

4. Clinical trials submitted to the journal as of July 1, 2018, must include a data sharing statement. The statement should indicate 1) whether individual deidentified participant data (including data dictionaries) will be shared; 2) what data in particular will be shared; 3) whether additional, related documents will be available (eg, study protocol, statistical analysis plan, etc.); 4) when the data will become available and for how long; and 5) by what access criteria data will be shared (including with whom, for what types of analyses, and by what mechanism). Responses to the five bullet points should be provided in a box at the end of the Methods section.

5. Standard obstetric and gynecology data definitions have been developed through the reVITALize initiative, which was convened by the American College of Obstetricians and Gynecologists and the members of the Women's Health Registry Alliance. Obstetrics & Gynecology has adopted the use of the reVITALize definitions. Please access the obstetric and gynecology data definitions at <https://www.acog.org/About-ACOG/ACOG-Departments/Patient-Safety-and-Quality-Improvement/reVITALize>. If use of the reVITALize definitions is problematic, please discuss this in your point-by-point response to this letter.

6. Because of space limitations, it is important that your revised manuscript adhere to the following length restrictions by manuscript type: Original Research reports should not exceed 22 typed, double-spaced pages (5,500 words). Stated page limits include all numbered pages in a manuscript (i.e., title page, précis, abstract, text, references, tables, boxes, figure legends, and print appendixes) but exclude references.

7. Specific rules govern the use of acknowledgments in the journal. Please note the following guidelines:

- * All financial support of the study must be acknowledged.
- * Any and all manuscript preparation assistance, including but not limited to topic development, data collection, analysis, writing, or editorial assistance, must be disclosed in the acknowledgments. Such acknowledgments must identify the entities that provided and paid for this assistance, whether directly or indirectly.
- * All persons who contributed to the work reported in the manuscript, but not sufficiently to be authors, must be acknowledged. Written permission must be obtained from all individuals named in the acknowledgments, as readers may infer their endorsement of the data and conclusions. Please note that your response in the journal's electronic author form verifies that permission has been obtained from all named persons.
- * If all or part of the paper was presented at the Annual Clinical and Scientific Meeting of the American College of Obstetricians and Gynecologists or at any other organizational meeting, that presentation should be noted (include the exact dates and location of the meeting).

8. The most common deficiency in revised manuscripts involves the abstract. Be sure there are no inconsistencies between the Abstract and the manuscript, and that the Abstract has a clear conclusion statement based on the results found in the

paper. Make sure that the abstract does not contain information that does not appear in the body text. If you submit a revision, please check the abstract carefully.

In addition, the abstract length should follow journal guidelines. The word limits for different article types are as follows: Original Research articles, 300 words. Please provide a word count.

9. Abstracts for all randomized, controlled trials should be structured according to the journal's standard format. The Methods section should include the primary outcome and sample size justification. The Results section should begin with the dates of enrollment to the study, a description of demographics, and the primary outcome analysis. Please review the sample abstract that is located online here: http://edmgr.ovid.com/ong/accounts/sampleabstract_RCT.pdf. Please edit your abstract as needed.

10. Only standard abbreviations and acronyms are allowed. A selected list is available online at <http://edmgr.ovid.com/ong/accounts/abbreviations.pdf>. Abbreviations and acronyms cannot be used in the title or précis. Abbreviations and acronyms must be spelled out the first time they are used in the abstract and again in the body of the manuscript.

11. The journal does not use the virgule symbol (/) in sentences with words. Please rephrase your text to avoid using "and/or," or similar constructions throughout the text. You may retain this symbol if you are using it to express data or a measurement.

12. Please review the journal's Table Checklist to make sure that your tables conform to journal style. The Table Checklist is available online here: http://edmgr.ovid.com/ong/accounts/table_checklist.pdf.

13. When you submit your revision, art saved in a digital format should accompany it. If your figure was created in Microsoft Word, Microsoft Excel, or Microsoft PowerPoint formats, please submit your original source file. Image files should not be copied and pasted into Microsoft Word or Microsoft PowerPoint.

Please upload each figure as a separate file to Editorial Manager (do not embed the figure in your manuscript file).

If the figures were created using a statistical program (eg, STATA, SPSS, SAS), please submit PDF or EPS files generated directly from the statistical program.

14. Authors whose manuscripts have been accepted for publication have the option to pay an article processing charge and publish open access. With this choice, articles are made freely available online immediately upon publication. An information sheet is available at <http://links.lww.com/LWW-ES/A48>. The cost for publishing an article as open access can be found at <http://edmgr.ovid.com/acd/accounts/ifaauth.htm>.

Please note that if your article is accepted, you will receive an email from the editorial office asking you to choose a publication route (traditional or open access). Please keep an eye out for that future email and be sure to respond to it promptly.

15. If you choose to revise your manuscript, please submit your revision via Editorial Manager for Obstetrics & Gynecology at <http://ong.editorialmanager.com>. It is essential that your cover letter list point-by-point the changes made in response to each criticism. Also, please save and submit your manuscript in a word processing format such as Microsoft Word.

If you submit a revision, we will assume that it has been developed in consultation with your co-authors and that each author has given approval to the final form of the revision.

Again, your paper will be maintained in active status for 21 days from the date of this letter. If we have not heard from you by May 03, 2019, we will assume you wish to withdraw the manuscript from further consideration.

Sincerely,

Nancy C. Chescheir, MD
Editor-in-Chief

2017 IMPACT FACTOR: 4.982
2017 IMPACT FACTOR RANKING: 5th out of 82 ob/gyn journals

In compliance with data protection regulations, you may request that we remove your personal registration details at any time. (Use the following URL: <https://www.editorialmanager.com/ong/login.asp?a=r>) Please contact the publication office if you have any questions.

To:
Nancy C. Chescheir, MD
Editor-in-Chief
Obstetrics & Gynecology
409 12th Street, SW
Washington, DC 20024-2188
Tel: 202-314-2317
Fax: 202-479-0830
E-mail: obgyn@greenjournal.org

May 2, 2019

Re: "Gabapentin for peri-operative pain management for uterine aspiration: a randomized controlled trial"

Dear Dr. Chescheir:

Thank you for your insightful comments on our manuscript. We appreciate the care that you and the reviewers have taken to evaluate our manuscript and are excited that you found it of interest. Your constructive comments on the overall theme of the paper, in regards to opioid use, as well as the presentation of the data are of great help. Below are point-by-point responses to all comments. We have revised our manuscript to reflect these concerns and look forward to your further comments. We are hopeful that these revisions make the manuscript appropriate for publication.

Best regards,

A handwritten signature in dark ink, reading "Jill Hagey". The signature is written in a cursive, flowing style. The first name "Jill" is written in a smaller, more compact script, while the last name "Hagey" is written in a larger, more expansive script with a prominent loop at the end.

Jill M. Hagey, MD MPH

On behalf of: Beverly A. Gray, MD; Donna Crabtree, PhD; Jeremy M. Weber, MB; Carl F. Pieper, DrPH; Lisa B. Haddad, MD, MS, MPH

REVIEWER COMMENTS:

Reviewer #1: This is an original research study describing a randomized controlled trial of adjunctive gabapentin for pain control during surgical abortion. The authors added 600mg gabapentin to their standard "minimal sedation" regimen and evaluated pain control during and at one day post procedure as well as anxiety, side effects and patient satisfaction. The authors found no difference in their primary outcome of pain at five minutes post procedure but did note significantly reduced opioid use post-procedure at one day. Importantly, the authors note widely varied practice in pain regimens post-surgical abortion and that gabapentin is a safe, well-tolerated medication that could reduce the need for opioid after a surgical abortion. Their study adds to the growing literature supporting use of gabapentin for abortion-related pain.

The study was IRB approved and registered on clinical [trials.gov](https://www.clinicaltrials.gov). The CONSORT checklist is included in the submission.

ABSTRACT:

1. Introduction: This is clear and defines up front what "usual" oral pain management means in your setting to set the context for readers.
 - a. Thank you for your thoughtful comment.
2. Methods: In line 38, I would add that you also measured patient satisfaction with pain control at one day post procedure.
 - a. While we did not measure patient satisfaction, we did measure patient pain perception at one day post procedure. We have revised the manuscript to reflect this measurement.
3. Results: In line 42-43, it would be important here to indicate that 73% of the participants received a narcotic prescription for post procedure pain control and that this wasn't standardized as part of the study protocol.
 - a. We have revised the manuscript to indicate the prescriptions of these narcotics were based on physician preference and not a part of the study protocol.
4. Conclusion: Line 45-46, this seems a bit vague to me here and in the discussion. I'm assuming this means studying gabapentin in local-only patients, or the effect on doses of IV opioids but if there is a way to make this more clear, that would be helpful to show the readers the implications of this study.
 - a. We have revised the manuscript to indicate that we were looking at the addition of gabapentin to our usual oral pain management regimens with paracervical block, and that we were specifically looking at oral opioid use one day after the procedure. Thank you for your request for clarification.
5. PRECIS: Need to add post-procedure opiate use since participants all still received pre-procedure opioid.
 - a. We have amended the PRECIS statement to clearly state that we are referring to post-procedure opioids.

INTRODUCTION:

6. In line 52, add in some examples of what else has been studied that you are referencing here. The generalist reader may not be familiar with this literature.
 - a. We have included information on other pain adjuncts as well as non-pharmacologic adjuncts to our introduction. We have also added an additional citation to provide more clarity to this overview of the literature.

7. Line 59, you start the introduction talking about surgical abortion in the US or North America but then reference uterine aspiration globally, do you mean internationally or globally like common? This is a really good point about using gabapentin for other similar gyn procedures but I was distracted by this.
 - a. We appreciate the reviewer's comment. We have changed this sentence to reflect that uterine aspiration is one of the most common surgical procedures around the world in addition to our discussion of practices in North America.
8. Line 64, I would indicate you also measured patient satisfaction with the experience and pain control
 - a. We have added that we measured pain perception at 24 hours after the procedure as detailed in the introduction. We have also deleted any comment regarding satisfaction in the manuscript to clarify this point.

MATERIALS AND METHODS:

9. Important that you included patients with chronic pain and/or history of drug alcohol use to make this result generalizable.
 - a. Thank you for your thoughtful comment.
10. Line 98, did you do any cervical prep for patients? If so how did that impact when/how they received their study med?
 - a. Patients above 12 weeks gestational age took a dose of vaginal misoprostol for cervical preparation 2 hours prior to their appointment. The timing of study medication was the same, which is now clarified in the methods section. As indicated in the "Drug Treatment and Procedures" section of the methods, all participants received the usual paracervical block with 40 ml of 0.5% lidocaine. We have clarified that the paracervical block is immediately prior to uterine aspiration, thus one to two hours following study medication administration.
11. Line 103, how long do patients in your clinic sit with lorazepam? Was the 1-2 hours for this study different?
 - a. Patients in our clinic outside of the study do typically wait at least one hour between receiving their usual perioperative pain medications and antibiotics prior to their procedure. There was no difference in wait time for the procedure based on this study. We have added the detail that this is standard practice in our clinic.
12. How did you decide on the five minutes after speculum removal pain score as the primary outcome? Is the standard time point the other pain studies cited use?

Previous studies have shown a variety of times for assessing pain, including immediately after the procedure or at the time of discharge from the procedure. On average most procedures last between 5 to 10 minutes in duration. In designing this trial, 5 minutes post-procedure was designated by a collective expert opinion.
13. Line 124, was there any standardization of the post-operative medications for the study or was it provider discretion? Do you have any providers who prescribe only NSAIDs post-procedure?
 - a. Post-operative medications were provided based on provider discretion, and were either ibuprofen or ibuprofen and oxycodone and acetaminophen. As noted in the abstract and results, 73% of women received a short-term prescription for oxycodone and acetaminophen, while the remainder 27% only received a prescription for NSAIDs post-procedure. We have added information regarding prescription of post-operative medications in the "Drug Treatment and Procedures" section.

RESULTS:

14. What percentage of your patients in this setting receive moderate IV sedation vs. oral pain management described in this study? Do you have patients that are local-only?
 - a. We do not have the capacity to provide IV sedation in the clinic setting and all patients receive oral sedation. Only a small percentage of our patients receive only local paracervical block (<5%) without adjunct opiate or anxiolytic medication.
15. Line 165, it is very impressive only one participant didn't complete the 24 hour phone call!
 - a. Thank you for the acknowledgement.
16. Can you provide the breakdown of indication for abortion by group either in Table 1 or in the paragraph lines 167-170?
 - a. Indication for the abortion was not collected as part of the participant survey, and is thus not reported as part of Table 1. Studies suggest that most women have more than one reason they obtain an abortion, making this data challenging to collect.
17. In line 193 on the influence of time, is this a result of the time they received the study med or the other oral sedation meds?
 - a. Thank you for your helpful comment. Participants received gabapentin or placebo at the same time as their other oral sedation medications, and so it is the time from administration of both that is being reported. We have clarified this within the "Influence of Time on Post-procedure Pain".
18. For women who received an opiate prescription, what was the dose and range of number given?
 - a. For women who received opiates, the dose was 5mg of oxycodone and 325mg of acetaminophen. The range of prescriptions given was between 5 to 20 tablets. This was based on provider preference and not regulated by the study protocol.
19. What were the results for patient satisfaction and were those different by group?
 - a. We did not measure patient satisfaction in this study. We did measure patient subjective self-report of pain at 24 hours post-procedure, which has been discussed in the "Post-operative Pain" section.

DISCUSSION:

20. Line 210 is too long for the first line of the discussion. The information in the second sentence should lead that paragraph.
 - a. We have revised the beginning of our discussion for clarity, focusing on our main findings in the first sentence.
21. Line 221 and line 223 seem to contradict each other. It is important to note that other studies have found that gabapentin impacted pain but what about uterine aspiration pain or the way you measured pain is different than those studies.
 - a. Thank you for your insightful comment. We have clarified these conflicting sentences to indicate the differences between our study and prior studies that have found that gabapentin impacted pain. We have included a recent study from JAMA Surgery that indicates that while gabapentin did not influence time to pain relief, it was associated with decreased opiate use, suggesting that it may take a longer period of time to see effects from gabapentin. We also noted that uterine aspiration pain was similar in each group.

22. Paragraph starting with line 234 is an important finding but needs background in the methods about how prescribing is done in your setting.
- Thank you for your helpful comment. We have added information in the methods and results section describing the opiate prescribing methods in our clinical setting. We also have clarified within the discussion that opiate prescribing was not significantly different between the gabapentin and placebo arms. Further, prescribers were blinded to treatment allocation, limiting any bias in different prescribing patterns.
23. The study in line 244 used pregabalin, this is an important distinction as pregabalin is much more expensive than gabapentin thus many sites have not adopted this evidence into practice.
- Thank you for your comment. While we agree that there are certainly practical differences between pregabalin and gabapentin, they are both well supported in decreasing peri-operative pain through a similar mechanism of action. We have noted that gabapentin was chosen in our study due to availability and cost, but we do believe we can compare our results to those presented here with pregabalin.
24. Is there a reference for line 249 regarding the variations in clinical practice around routine prescribing of opioids post abortion?
- Thank you for your question regarding routine prescribing of opioids post abortion. In a survey of National Abortion Federation members performing first trimester abortion procedures, approximately 30% of facilities either prescribed or gave tablets of opioids following the procedure. We have included this citation in our manuscript. As mentioned, prescription of opioids post-abortion was left to provider discretion in our study.
25. A strength I would include is that you measured baseline pain, history of opiate use and sexual trauma.
- Thank you for your comment. We have included this as a strength.
26. In line 259, that data isn't presented in your results, readers can determine generalizability based on the demographics in Table 1 and how similar the protocol for oral sedation followed is to their clinical practice. While similar between study arms, having 31% of participants with a master's degree and reporting a household income of \$>100,000 should probably be remarked on in the discussion and reflect the setting the study was done in perhaps.
- We have revised our discussion to reflect the possibility of skewed demographics in this study.
27. The last paragraph should be strengthened, I would be clear in the finding that gabapentin wasn't associated with decreased pain at 5 minutes after oral sedation for first trimester aspiration abortion but was well-tolerated and lead to reduced opiate use post-procedure which is what this study adds to the current literature. If you have recommendations for further research or plan to evaluate this relationship further, add that in too.
- Thank you for your comments. We have re-worked the conclusion to more clearly state our primary outcome. Additionally, we have added more information on what further studies would advance the field including specifically the utility of gabapentin if given at other timepoints, such as one day prior to the procedure at the time of laminaria placement for second trimester procedures, its use among select populations, or as an adjunct to other pain management strategies such as IV sedation or paracervical block alone.

TABLES AND FIGURES

28. Table 1, can you add in indication for abortion?
 - a. **As in the body of the text, we did not collect the indication for abortion.**
29. Table 4, can you add in nausea/vomiting outcomes?
 - a. **Nausea and vomiting have been added to Table 4 as noted, with no statistically significant difference in nausea VAS score or vomiting at 10 minutes post-procedure.**

Reviewer #2:

TITLE: Gabapentin for uterine aspiration

Overall: This is a randomized controlled trial comparing the addition of gabapentin to placebo for pain management during and after uterine aspiration procedures for induced abortion. 96 women were enrolled. The study was conducted at a single institution.

Overall in the abstract, and elsewhere in the paper there is an jump from gabapentin related pain score and opioid use. This jump is implicit and not an actual measurement. Please clarify the measures in this study - VAS and gabapentin or opioid use.

Many physicians do not use opioids at all for post-op pain management after first trimester uterine evacuations so please be clear when you use it and when you don't. This paper should not ultimately serve as justification for increased use of post-operative opioids for a very minor procedure.

Does abstract provides good information but there are some places where the abstract could be stronger:

1. The objective written in the abstract should be the same as the objective and primary outcome - which is pain 5 minutes post procedure.
 - a. **We have amended the abstract to have the objective reflect the primary outcome.**
2. For participant flow please state the overall number screened, enrolled and included in the analysis groups per the CONSORT guidelines recommendation. Providing the screened number matches your opening statement in the results which includes the statement about the women who were excluded from enrolling.
 - a. **We have included the number of patients screened in the abstract.**
3. Overall in the abstract, and elsewhere in the paper there is an jump from gabapentin related pain score and opioid use. This jump is implicit and not an actual measurement.
 - a. **We appreciate the reviewer's comments. The purpose of this manuscript is twofold. First, we report the primary outcome of a randomized controlled trial on the effect of peri-procedural gabapentin prescription on pain five minutes post procedure. As the reviewers have pointed out, this outcome was not significantly different between groups. Additionally, we exhibit data that women who did receive gabapentin periprocedurally utilized less opioids in the days following the procedure. We have revised the general tone of our manuscript in multiple places to more clearly delineate between the two findings, more definitively state that this was a negative study, and more explicitly define our finding of decreased opioid use as a secondary outcome. While secondary, we believe that decreased opioid use is of interest to the gynecologic community as there are a significant**

number of abortion providers who do indeed prescribe opiates at the time of first trimester aspiration. As excessive opioid use is of great contemporary concern, we believe it is appropriate to highlight this finding in our study specifically and advocate for more uptake of and research around interventions periprocudurally to decrease post-procedure pain.

How this paper fits in with existing literature:

Author Conflicts are stated. The paper will be presented at ACOG annual meeting in May.

IRB/Ethics approval was obtained from Duke IRB

If RCT was it registered? Yes, First Posted: April 1, 2016; Last Update: November 8, 2018, as completed. Primary outcome listed as pain score on 11-point VAS 5 minutes post procedure.

The outcomes measures and intended sample on Clinical Trials site agree with those in paper.

If the CONSORT guidelines were used please state such and reference them in the paper.

CONSORT Statement: David Moher, Sally Hopewell, Kenneth F. Schulz, Victor Montori, Peter C. Gøtzsche, P.J. Devereaux, Diana Elbourne, Matthias Egger, Douglas G. Altman, CONSORT 2010 Explanation and Elaboration: updated guidelines for reporting parallel group randomised trials, Journal of Clinical Epidemiology, Volume 63, Issue 8, August 2010, Pages e1-e37, ISSN 0895-4356, <http://dx.doi.org/10.1016/j.jclinepi.2010.03.004>.
(<http://www.sciencedirect.com/science/article/pii/S0895435610001034>)

TITLE: the title is good - succinct and meets CONSORT guidelines.

PRECIS: State the precis in active language.

- a. Thank you for your comments, we have changed the PRECIS to be in the active voice.

INTRODUCTION:

1. The introduction could be a little more specific about the procedure (first trimester vacuum aspiration) and usual pain management and then the role of this with the opioid epidemic - if it fits. Overall the references to the opioid epidemic seem a bit of a tangent.
 - a. We have revised our abstract to include the exact procedure studied. As discussed above, we have clarified our language regarding opioids and our primary and secondary outcomes.
2. Lines 58: Please include in the introduction an overview or statement of how frequently opioids are actually used for postoperative pain management after first trimester uterine evacuations. Certainly in this reviewers
 - a. Thank you for your suggestion. As mentioned above, approximately 30% of facilities either prescribed or gave tablets of opioids following first trimester abortion procedures. We have included this citation in our manuscript.
3. Need to be specific - uterine aspiration or dilation and evacuation.
 - a. Thank you for your helpful comment. We have revised the language to specifically state that we used manual vacuum aspiration or electric uterine aspiration as opposed to dilation and curettage, both in the introduction as well as the methods. Patients under 11 weeks gestational age underwent a manual vacuum aspiration, while those at 11 weeks gestation or above underwent an electric uterine aspiration.
4. Line 62: Please be clear - if you assessed please be clear which is the primary objective and which is secondary. The objective should match the outcomes and the reporting.

- a. We have revised this section to clearly state that our primary outcome was pain five minutes post-procedure and our secondary outcomes were anxiety, side effects, and post-operative pain medication use.

MATERIALS AND METHODS:

5. Overall the methods could be more concise.
 - a. Thank you for your comment. We have shortened the methods section for clarity.
6. The section on outcomes should be concise and focus on the primary and secondary outcomes. Currently there is so much information in the paragraph the details and focus is hard to follow.
 - a. We have revised this paragraph to focus on our primary and secondary outcomes. We have also made this section more concise per the reviewer's request.
7. Lines 92-94: Please state who prepared the blinding for the study medications.
 - a. Our investigational drug service pharmacists prepared blinding for the study medication. We have revised the manuscript to describe this procedure.
8. Line 97: Headings for sections aren't needed if strong enough topic sentences are included.
 - a. Thank you for your comment. Given that this study had multiple different assessments, we believe that these headings help provide clarity for the reader.
9. An overall statement of the type of study should be made at the beginning of the Materials and Methods section - cross-sectional? Retrospective or prospective cohort?
 - a. Thank you for your suggestion. We have clarified in our first sentence of the Methods section that this was a prospective, randomized, double-blind, placebo controlled trial.
10. Is the population and setting clearly stated? The population and setting should be clearly stated - what are the demographics of patients and clinicians at the setting the study took place?
 - a. We appreciate the reviewer's comment. We have included more information about the Duke Family Planning Clinic and the clinical staff of the clinic.
11. The subject selection is clearly stated? concise with a list of eligibility criteria only - this present the combination of positives and negatives being listed.
 - a. As stated in the manuscript, we have clearly outlined both the inclusion and exclusion criteria, as well as how participants were assessed for eligibility.
12. Does the process minimize bias?
 - a. As stated in the manuscript, we minimized bias by randomizing participants to gabapentin versus placebo and by blinding both participants and providers to study group allocation.
13. The primary outcome is clearly stated:
 - a. As stated in the manuscript, the primary outcome was pain five minutes after removal of the speculum measured on a 100-mm VAS score.
14. Adverse events appropriately collected?
 - a. As stated in the manuscript, adverse events were collected by a checklist of side effects that was assessed prior to discharge.
15. Is sample size justified? For which outcome?
 - a. Our sample size was justified for our primary outcome of pain five minutes after removal of the speculum on the 100-mm VAS based on an assumed standard deviation and a power to detect a 15-mm difference with a power of 80% and a type 1 error rate of 0.05.

RESULTS:

16. A participant flow chart should be included - this will provide readers with an understanding of the final analysis population, and this is also a crucial component of reporting according to STROBE.
 - a. We have included a participant flow chart as Figure 1 of our paper using the CONSORT flow diagram. As we are reporting a randomized, controlled trial, we believe that the CONSORT flow diagram is more appropriate for our participant flow chart than the STROBE flow diagram.
17. Please include a statement of how many women were approached/screened/enrolled in the opening of the results.
 - a. As stated in the manuscript, we approached 113 women for the study, after which 96 were screened and enrolled in the study.
18. Is participant flow chart included?
 - a. As above, we have included a participant flow chart as Figure 1.
19. Response rate/follow-up rate good enough?
 - a. We believe our response rate and follow-up were excellent with only one participant not receiving their allocated intervention and only one participant lost to follow up at the 24 hour post-procedure phone call. This is included in the manuscript.
20. Is bias minimized? Is confounding considered?
 - a. As above, we minimized bias through randomization and double-blinding as described in the methods section. As our randomization was successful in obtaining similar demographic characteristics between groups, we did not control for any additional confounders.
21. How are dropouts defined and analyzed?
 - a. Dropouts were defined as participants who did not receive their allocated intervention. As this participant dropped out prior to any intervention, they were not included in an intention to treat analysis. This information is included in the manuscript.

DISCUSSION:

22. Please state the overall findings of your study in the context of your population and setting - this study is limited in that it's single institution.
 - a. We have revised the manuscript to more clearly state that this was a single institution study and also to highlight how the characteristics of our study sample may not be broadly generalizable.
23. The fact that MORE opiate was used by gabapentin group than placebo group actually demonstrates that there may be some harm when gabapentin is used - this must be reported and addressed in the discussion.
 - a. Among participants prescribed opiates in the post-operative period, there was a non-significant trend towards more opiate prescriptions in the gabapentin group (80.9% versus 64.6%). We are commenting on the actual use of these opiates in the post-operative period, where participants in the gabapentin arm used significantly less opiates than those in the placebo group (18% versus 47%, $p = 0.01$). We have revised this section for clarity.
24. Overall the discussion could be a bit shorter.
 - a. The discussion has been shortened for clarity.
25. Lines 234-243 - this is important and is related to comments about the overall paper - implying that gabapentin may reduce opiate use...I think this just detracts from the

overall paper. What was the rationale for the study? To decrease opiate use or to decrease pain associated with uterine aspiration in first trimester for induced abortion?

- a. Thank you for your comments. As we have stated above, the intent of the randomized controlled trial was to decrease peri-procedural pain. However, an interesting finding is that gabapentin decreased later opioid use. We have revised the manuscript in multiple ways to make this more clear.
26. Are limitations discussed? Are any insurmountable?
- a. Limitations to the study include assessment of pain at only a few time points and that this was a single clinic trial, as noted in the discussion.
27. Is conclusion consistent with results?
- a. We believe the conclusion that gabapentin was not associated with decreased pain at five minutes post procedure to be consistent with our results and representative of our primary outcome.

TABLES AND FIGURES

28. The tables should have more descriptive figures. The tables would be easier to read esp Table 1 if the percentage signs were removed, numbers were lined up, and percentages were rounded.
- a. We have revised the table for readability by removing percentage signs, ensuring correct alignment and rounding to the nearest percent.
29. There should be a table that clearly demonstrates opiate use during and after the procedure by group since this was a secondary outcome, and a lot of text is devoted to this concept.
- a. Thank you for the helpful comment. We have added Table 5 to indicate the use of post-operative pain medication use by gabapentin versus placebo group to present the data related to opioid use.

Reviewer #3: Thank you for your submission.

Please see comments below:

1. Appreciate the inclusion of women with chronic pain in the study as excluding them would have decreased the ability to generalize the results.
 - a. Thank you for comments. We agree that it was important to include women with chronic pain.
2. Would appreciate more information regarding what the "perioperative clearance" entailed
 - a. We have included more information on perioperative clearance. Essentially, this entailed assessing patients for an in clinic procedure. We take care of many medically complex patients, some who require more monitoring or greater sedation than oral sedation. Patients are also offered a procedure in the OR if they desire.
3. The 24hr pain scale used was the reverse of the traditional pain scale used in across medical specialties (where 0=no pain and 10=most severe) which could potentially confuse patients (or even data collectors).
 - a. Thank you for your comments. We agree that the pain scale is the reverse of the traditional pain scale. However, this pain scale is based on the quality of recovery survey, which is a validated survey that has been extensively studied in the post-

operative period among many different patient populations. Information about this pain scale is highlighted in the results section.

STATISTICAL EDITOR COMMENTS:

The Statistical Editor makes the following points that need to be addressed:

1. lines 112-113, 132-136: The primary outcome was specifically cited as VAS on a 100-mm scale at 5 minutes after speculum removal. The primary outcome should first be reported in the same format. That is, as two VAS 100 point VAS scales for the gabapentin vs control cohorts and cited as means(SD) with stats testing. If the distributions were more appropriately non-parametrically tested as median(Range or IQR), that could then be cited, but not as the primary outcome. Likewise, the correction for the baseline pain scale, although of interest, was not the primary outcome. Furthermore, the cohorts were randomized, so any difference in baseline pain scores would be attributable to random chance. That is, the primary outcome should be clearly demarcated from the other, secondary outcomes
 - a. Thank you for your comments. We have revised the manuscript to be more focused on our primary outcome and more clearly demarcated the primary outcome from the secondary outcomes. We acknowledge and understand your comment on correcting for baseline pain. As cited in the manuscript and again below¹, we decided to correct for baseline pain for multiple reasons.
 - i. To reduce any residual baseline differences between the treatment arms
 - ii. Controlling for baseline pain reduces the variance, which in turn yields narrower confidence intervals and a more precise estimate for the treatment effect
 - iii. The best predictor for a future outcome is an earlier measurement of that same outcomeTo be clear, baseline pain was not post-hoc adjusted for due to observed differences in that variable between treatment arms. The decision was pre-specified in the statistical analysis plan knowing baseline pain and pain 5 minutes after the procedure would be associated. As a sensitivity analysis, the primary outcome was compared between treatment arms using a t-test. The treatment effect moved slightly closer to the null and remained insignificant. Reporting of the primary outcome through the manuscript has been changed from median (Q1, Q3) to mean $\pm$ SD. In the manuscript, the primary outcome was tested parametrically using linear regression. The normality of the residuals was checked and the assumption looked plausible.
 1. Committee for Proprietary Medicinal Products (CPMP). Points to Consider on Adjustment for Baseline Covariates. Stat Med 2004;23(5):701-9.
2. General: The precis, abstract and results sections need to be re-written, so as to emphasize that the primary outcome of this RCT was negative. Some secondary outcomes were statistically significant, but those were not how the hypothesis was stated nor how the power/sample size calculated.
 - a. As above, we have revised the manuscript to focus more on the primary outcome.
3. Table 1: Need units for age, income (annual in dollars)
 - a. We have included units for age and income in Table 1.

4. Table 3: Were the anxiety scores normally distributed? If not, should cite as median(range or IQR) and test non-parametrically.
 - a. These data are not normally distributed. Anxiety scores have been changed from mean (SD) to median (Q1, Q3). Wilcoxon rank sum test was used to compare anxiety scores between treatment arms.

EDITOR COMMENTS:

1. Thank you for your submission to Obstetrics & Gynecology. In addition to the comments from the reviewers above, you are being sent a notated PDF that contains the Editor's specific comments. Please review and consider the comments in this file prior to submitting your revised manuscript. These comments should be included in your point-by-point response cover letter.

The notated PDF is uploaded to this submission's record in Editorial Manager. If you cannot locate the file, contact Randi Zung and she will send it by email - rzung@greenjournal.org.

1. My major concern for you to address in your revision is that this is essentially a negative study. For your primary outcome, there was no difference between gabapentin and placebo. Your paper as written sort of glosses over that and instead focuses on the issue of side effects and post op opioid use, which were secondary outcomes. The use of opioids post these procedures should be quite low to begin with in the average patient. The focus on the positive secondary outcomes, while not focusing on the negative primary outcome is a form of what is known as "Spin" and needs to be corrected to present this as a negative outcome, with the secondary, exploratory outcomes worth studying further.
 - a. Thank you so much for your thoughtful comments. We have revised our manuscript to indicate that this was a null study, and our findings regarding opioid use were part of our secondary analysis. As indicated in the reviewer comments,
2. - The Journal style doesn't not use the virgule (/) except in numeric expressions. Please edit here and in all instances.
 - a. We have removed the virgule in all inappropriate instances.
3. - In both the abstract and the paper, please provide absolute numbers as well as which ever effect size you are reporting (if appropriate) + Confidence intervals. P values may be omitted for space concerns. We strongly prefer CI's as they give more information about strength of association than do P values. By absolute values, I mean something like xx (outcome in exposed)/yy(outcome in unexposed) (zz%) (Effect size= ; 95% CI=.) An example might be: Outcome 1 was more common in the exposed than the unexposed 60%/20% (Effect size=3;95% CI 2.6-3.4)
 - a. Thank you for your feedback. We have revised the text in both the abstract and the paper to report absolute numbers, an effect size, and confidence interval for all parametrically tested outcomes.
4. - When you write that a study occurred between date 1 and date 2, it literally excludes those boundary dates. For instance, "This study was performed between Feb 2018 and Jan 2019" would mean it was performed from March 2018 to Dec 2018. Do you instead mean that the study was performed from date 1 to date 2? If so, please edit. The same is true on line 32 in description of the included EGA's.
 - a. We have revised dates in the manuscript to reflect more clearly the actual time during which the study was performed.
5. - plural: procedures

- a. Thank you for your comments, we have made this plural.
6. - Journal style uses cesarean birth or delivery but not cesarean section.
 - a. We have revised all instances of "cesarean section" to read "cesarean birth".
7. - It is an idiosyncratic fact that at the Journal we tend to avoid the use of the word impact to imply the result of a change, preferring to limit "impact" to mean a physical blow.
 - a. We have removed the word impact from the manuscript.
8. - For clarity, in some places you state inclusion was of patients SCHEDULED to undergo surgical abortion and here you say it was presentation. As these are potentially different #'s of women, which is it? Were the randomized at time of scheduling or at presentation?
 - a. We have clarified this in the methods section to indicate that women were approached at the time they arrived for their scheduled procedure appointment.
9. - These semi-colons should be commas.
 - a. We have revised this punctuation.
10. - who approached them?
 - a. Study subjects were approached by either the clinic nurse or physician. This has been revised in the manuscript.
11. - is this on the day of surgery?
 - a. Thank you for your comment. This was on the day of surgery, which is better reflected in the methods section.
12. - Could you combine the information on lines 106-108 and 113-116?
 - a. We have combined the information on lines 106-108 and 113-116.
13. - You were doing 3 different VAS measurements at each of these time points on a patient w/ narcotics and a anxiolytic on board. This reasonably could have taken more than 1-2 minutes to perform. Did you standardize the order of administration? Since it was. your primary end point, was the 5 minute post speculum removal pain score done first, before the anxiety and nausea scores?
 - a. All surveys were administered by the study team during the procedure. Patients were very adept at designating their pain/anxiety scores on the VAS scale, and it would usually take <30 seconds to make the 3 separate marks on each scale, not adding any additional time to the procedure. Pain was assessed first, followed by anxiety and nausea for all participants.
14. - what is your standard prescription for women to go home with for pain management? Did it differ by EGA?
 - a. The standard prescription was determined based on provider. It is generally 800mg ibuprofen with or without oxycodone. We have included more detail regarding these prescribing practices in our manuscript.
15. - For an RCT, is is unusual to have exactly equal numbers
 - a. Thank you for your comment. Though unusual we were able to achieve this because of our block randomization scheme that had equal numbers in each group. As our randomization occurred just prior to medication administration, patients were very unlikely to discontinue their study participation at that time.
16. - please state this the way you do in methods "five minutes after removal of speculum" to avoid confusion,
 - a. We have added "five minutes after removal of speculum" for clarity.
17. - are these different? Please provide statistical information as noted above in the abstract comments
 - a. We have added this statistical information as noted in the abstract comments.
18. - We do not allow authors to describe variables or outcomes in terms that imply a difference (such as of the terms "trend" or "tendency" or "marginally different") unless there is a statistical difference. Please edit here and throughout. Here you should start the sentence by saying "Pain five minutes after removal of the speculum, controlling for

baseline pain, was not statistically different from the nul. (xxxx) with only a 3.4 mm difference. Same true for sentence on line 176

- a. We have revised all instances of implied difference to reflect the fact that there was no statistically significant difference.

19. - Are these reversed? This is opposite of the design of a normal pain scale. Could that have been a source of confusion for patients?

- a. Thank you for your comment. As detailed above, the pain scale for the VAS is opposite of the pain scale for the moderate and severe pain in the quality of recovery survey. This pain scale is based on the quality of recovery survey, which is a validated survey that has been extensively studied in the post-operative period among many different patient populations. As this survey was completed at a different time point than the VAS pain scales (24 hours post-procedure versus immediately during and after the procedure), there was less likely to be confusion between the two pain measurements.

20. - please provide the absolute numbers as well.

- a. Thank you for your helpful comment. The absolute numbers have been included.

21. - Please be very careful about using "after the procedure" as it is a bit vague. Line 181 you talk about "on removal of the speculum and end of the procedure"--I assumed those were 2 different times. Here on line 189 you say "5 minutes after the procedure"...was that at speculum removal or some other time.

- a. We have clarified the timing in all instances to specifically state exactly when a measure was taken.

22. - was this the same for both groups?

- a. It was the same for both groups, and we have clarified this point in the manuscript.

23. - hope in discussion you comment on ability to decrease prescription for opioids overall, independent of whether you continue to use gabapentine.

- a. Thank you for your constructive comment. We have added this point to our discussion.

24. - This is called a primacy claim: yours is the first, biggest, was innovative, etc...In order to assert that, you need to provide the search terms used and the data base (s) searched (PubMed, Google Scholar, etc) to substantiate the claim. Otherwise, it needs to be deleted.

- a. We have deleted this primacy claim.

25. - I thought you had one who did not follow up at 24 hours? (Line 165)

- a. There was indeed one patient who did not participate and therefore did not follow up. This is corrected for clarity of this point.

26. - excellent graphic

- a. Thank you for your comment regarding our graphic. We believe it is helpful in illustrating our point from the manuscript.