OBSTETRICS & GYNECOLOGY

NOTICE: This document contains correspondence generated during peer review and subsequent revisions but before transmittal to production for composition and copyediting:

- Comments from the reviewers and editors (email to author requesting revisions)
- Response from the author (cover letter submitted with revised manuscript)*

**The corresponding author has opted to make this information publicly available.*

Personal or nonessential information may be redacted at the editor's discretion.

Questions about these materials may be directed to the *Obstetrics & Gynecology* editorial office:
obgyn@greenjournal.org.

Date: Jul 25, 2019
To: "David Sheyn" [REDACTED]
From: "The Green Journal" em@greenjournal.org
Subject: Your Submission ONG-19-1161

RE: Manuscript Number ONG-19-1161

Development and Validation of a Machine Learning Algorithm for Predicting Response to Anticholinergic Medications for OAB Syndrome

Dear Dr. Sheyn:

Your manuscript has been reviewed by the Editorial Board and by special expert referees. Although it is judged not acceptable for publication in Obstetrics & Gynecology in its present form, we would be willing to give further consideration to a revised version.

If you wish to consider revising your manuscript, you will first need to study carefully the enclosed reports submitted by the referees and editors. Each point raised requires a response, by either revising your manuscript or making a clear and convincing argument as to why no revision is needed. To facilitate our review, we prefer that the cover letter include the comments made by the reviewers and the editor followed by your response. The revised manuscript should indicate the position of all changes made. We suggest that you use the "track changes" feature in your word processing software to do so (rather than strikethrough or underline formatting).

Your paper will be maintained in active status for 21 days from the date of this letter. If we have not heard from you by Aug 15, 2019, we will assume you wish to withdraw the manuscript from further consideration.

REVIEWER COMMENTS:

Reviewer #1:

Summary: This is a very interesting article that addresses an important clinical question. Often times we treat OAB patients with multiple medications before moving on to 3rd line therapy. With this model, we could potentially triage patients faster and help counsel patients on their risk of treatment failure.

Abstract:

*Line 62, delete the "." after practice

Methods:

- * Can more information be given on the TRIPOD statement .
- * Were patients excluded if they had tried an OAB med prior to coming to your office?
- * How was treatment response of > 50% documented.. was this the stand at your institution? Was a validated questionnaire used?

Results:

- * Line 270: change "never" to no
- * What was the distribution for the other medications prescribed?
- * If the majority were treated with oxybutynin or tolterodine, is your model accurate for all the anticholinergic medications?

Conclusion:

- * Line 321: delete "are" before generally
- * Additional limitation is that all data came from single institution with predominately white and African American population.
- * I would also add that no validated questionnaires were used. Most studies these days have a composite score for success, both subjective and objective. This has been shown to have higher correlate with success.

Reviewer #2:

1. The objective was clearly defined and relevant to literature for future personalized medicine in treatment of OAB
2. Line 161 discussing follow up: were patients who never reached maximum dosing excluded and if so how do you feel this would impact prediction of all patients starting on treatments for OAB
3. Explain why only ER medication prescriptions were only used for the study inclusion and search-again this could increase the number of inclusion in data analysis and more generalizable data
4. Tables need titled and numbered. Difficult to read Not certain of usefulness to readers of having the Random forests noted in the journal
5. Overall commend authors on use of excellent data set and relevant inclusion. Other factors to consider would be pad use, use of depends, and if possible a validated questionnaire at baseline such as MESA scores since this is very subjective and the study in terms of definition of success and failures are also subjective this could lead to understanding if high bothersome is significant in predictive model to failure
6. Results: would recommend limited some of the repeat data being reported from Table 3
7. In Discussion: limitation of study should include short follow up median 97 days in data. Clinically when counseling patients, longer term follow up in success would be more relevant -This would not help predict the <20% who continue to use AC after one year which would be more clinically relevant.
8. Conclusion- again accurate prediction in average follow up of 3 months should be stated

Reviewer #3:

Abstract - Objective: To develop and externally validate a prediction model for anti-cholinergic response in patient with Overactive Bladder (OAB)

Methods - model retrospective dataset of 559 women with OAB treated with anti-cholinergics between 1/2010-12/17 primary outcome was >50% improvement of patient reported symptoms, groups stratified by age <40, 40-60, >60 and failed AC medications

stratified cross validation with machine learning algorithms - Random Forest (RF) was most accurate treatment failures or responders; RF model externally validated - prospective collected data in 82 women

Results - correctly predicts failure in 81% and response in 61%

Conclusions - validated algorithm can predict AC failure

Intro - OAB incidence, treatment is behavioral and pelvic PT

AC tx - average efficacy 50-60% with <20% continuing after 1 year, goal is to determine which patient benefits from prediction model

objective: develop and validate predictive model to determine likelihood of success or failure of AC treatment using machine learning methods

Materials/ Methods - IRB approval, use of multivariable prediction model for treatment of OAB with AC medication from 1/10-12/17

Patient selection - identified by prescriptions in medical records, inclusion and exclusion criteria are listed data for 41 clinical risk factors collected with primary outcome being subjective failure defined as no or minimal improvement after a minimum of 4 weeks of treatment on maximum dose with documentation of medication adherence

Model development - patient stratified by age and previous failures

8 features for model, ROC threshold defined, patients classified as responders and non responders

External validation - data on same parameters collected 1/17-12/18 and accuracy/ AUC assessed

Results - 559 patients were followed for 97 days - 47.6% with treatment response

accuracy of RF model 80.3% - most predictive for patients <40 and least predictive for patients 41-60 with previous failure External validation - 82 in external validation, sensitivity and specificity of model with treatment failure as outcome 79.2 and 79.3%

most predictive for patients <40 and least predictive for >60 with previous failure

37.6% responded to treatment and those who didn't respond were correctly predicted in 81%

Discussion - AC medications have a modest improvement, machine learning algorithm uses model to discriminate likelihood of treatment failure

good accuracy in women <40 and >60 if they haven't previously failed

strengths - large sample size with external validation

limitations - retrospective, subjective

developed and validated a model - prospective trials with voiding diaries are needed

Comments:

- 1) Entire manuscript needs to be thoroughly proofread for multiple errors - grammatical, syntax, and spelling- throughout but particularly in the results and discussion part
- 2) 209-218 - clarify and proofread model technique - please make more comprehensible
- 3) 240-243 - proofread
- 4) why is the model prediction different for the cases than the external validation - what does this reflect? results - most predictive for patients <40 and least predictive for 41-60 with previous failure but external validation most predictive in <40 and least predictive if >60 with previous failure
- 5) clarify what a machine learning model is and how it factors in the variables, which variables have which impact. It would be helpful to know which factors put people at risk for failure
- 6) The idea of a model is interesting, but if only 37% of the people respond, please indicate what added value this provides - it is already clear that most people won't respond regardless and should be counseled as such
- 7) are these patients also pursuing pelvic PT - if so, this could confuse the results/responses

STATISTICAL EDITOR COMMENTS:

The Statistical Editor makes the following points that need to be addressed:

General: In my version of the article, the Tables and Figures are not labelled, so I will cite them by their numbers given in the text, with, with a short description, as needed.

General and lines 191-198: Since this is not a method our readers would be accustomed to, the Authors should explain how a sufficient sample size vs the proportion of adverse outcomes and how to a priori limit the number of features in the model. That is, how was the number of features in this case chosen as 8? Would a more parsimonious model have similar accuracy and performance? How is the trade-off ascertained between adding more features vs incremental gain in accuracy?

lines 52-53: Need to clearly identify how many women in the training set and the validation set successfully met the definition of the primary outcome, that is, how many treatment failures or responders there were overall, and also within the various subsets.

lines 66-71: See later comments re: CIs, for both the training and the validation set.

Table 3 (Training group characteristics): I presume the hemoglobin A1c were only from the 124 women with DM, but should so state, Should also enumerate all missing values. The last entry "Mean time to first follow-up ...) is formatted as median(IQR), so need to clarify that 97 days represents a median, not a mean value.

Table 3 and lines 204-207 (list of RF, SVM and tree augmented Naive Bayes vs 5 combinations of age and NOF): This list of accuracies needs to include CIs, based on the sample sizes. One cannot state that one method is superior because the absolute value of the accuracy is highest. Need to compare statistically, since N = 559 did not represent all women with OAB, but was a sample.

lines 280-283: Similarly, the AUCs all need CIs, based on the sample sizes for their subsets by age, and "most predictive" must be based on stats comparisons, not simply which one had the highest arithmetic proportion.

lines 298-301: Need to include CIs for the sensitivity and specificity. Again, the comparison of AUCs needs CIs and stats testing. I presume this paragraph is referring to the external validation set only, since that was the subheading on line 285. If this refers to the larger data set, or some combination, should insert a different subheading on line 297.

lines 303-306: This section's arithmetic seems inconsistent with the total number of women in the test set and internally inconsistent. For example, since the N = 82, then if 31 responded, then the response rate would be 37.8%, not 37.6%. Assuming that 31 responded, then 51 did not, and the model would correctly predict 81.2% of those, which is 41.4. If the overall treatment response was correct 61.5% of the time, that would be 50.4 women. Something is not right. Need to cite internally consistent numbers and CIs. If in fact the correct response of ~ 61% had a lower bound CI of 50%, then the model was statistically no better than a coin toss.

Table 5, lines 286-287: Same issue with "Mean time ..." of mean vs medians. Need to enumerate all missing values and

clarify that the hb A1c were only from the DM subsets.

The figures showing decision trees need further explanation. No explanation is given for the gini coefficient, and most of our readers will not be familiar with it, for example. Might be worthwhile to focus on one section of the tree and explain in more detail what is represented.

The figures of AUCs need figure legends and stats corroboration of any statement of superiority. Most of the AUC do not clearly separate, often cross one another and there is no clear hierarchy, except possibly for the highest vs the lowest.

EDITOR COMMENTS:

1. When you review your paper, I recommend that you read it aloud at some point. Throughout your first submission, there are a number of missing article and pronouns, and incorrect tense use. I've pointed out some of these, but it will perhaps help if you read the paper.

2. Thank you for your submission to Obstetrics & Gynecology. In addition to the comments from the reviewers above, you are being sent a notated PDF that contains the Editor's specific comments. Please review and consider the comments in this file prior to submitting your revised manuscript. These comments should be included in your point-by-point response cover letter.

The notated PDF is uploaded to this submission's record in Editorial Manager. If you cannot locate the file, contact Randi Zung and she will send it by email - rzung@greenjournal.org.

- You will need to include a précis: The précis is a single sentence of no more than 25 words, written in the present tense and stating the conclusion(s) of the report (ie, the bottom line). The précis should be similar to the abstracts conclusion. Do not use commercial names, abbreviations, or acronyms in the précis. Précis should be the "hook" for people who scan the Table of Contents to see what to read. It shouldn't not include statements like "in this study" or "we found". Just state what you found. It may be that the conclusion of your abstract as edited will be good.

- We no longer require that authors adhere to the Green Journal format with the first submission of their papers. However, any revisions must do so. I strongly encourage you to read the instructions for authors (the general bits as well as those specific to the feature-type you are submitting). The instructions provide guidance regarding formatting, word and reference limits, authorship issues, and other things. Adherence to these requirements with your revision will avoid delays during the revision process, as well as avoid re-revisions on your part in order to comply with the formatting.

- The way this is worded, I'm unclear if your paper is a secondary analysis of another study or not. I think what you mean is "For the model development, the primary outcome was a >50%...." . Is that correct?

- what is NOF?

- please be consistent. Here you use AUROC; line 66 you use AUC, which is the more common.

- I'm confused why you used treatment success in the model development phase of your study but treatment failure in the validation portion. Why not be consistent?

- likelihood ratio of what?

- about 80% of the time.

- Although its not required by the Journal, you. may wish to reference and use the publication guideline for machine learning predictive models. They describe K folding quite well and this may be a useful reference in that regard.

J Med Internet Res. 2016 Dec; 18(12): e323.

Published online 2016 Dec 16. doi: 10.2196/jmir.5870

PMCID: PMC5238707

PMID: 27986644

Guidelines for Developing and Reporting Machine Learning Predictive Models in Biomedical Research: A Multidisciplinary View.

- For parsimonious word choice would you consider: "In patients who fail to respond to these treatments, pharmacotherapy is the next option with anti-cholinergic medications the most common choice. "

- what is 3rd line therapy if not medications? I'm a bit confused by lines lines 123-125. For the 40-50% of women who do not respond to AC treatment, are there other pharmacotherapeutic options? Is it common to cycle through the several different AC meds before calling the drug class a failure?

- validate A predictive model
- name the IRB here
- Not sure what you mean by this phrase.
- This is the EMR?
- What if patients were tried on several of these drugs?
- For clarity here is it with or without frequency AND with or without urgency AND with or without Nocturia?
OR is it urgency with or without frequency, OR incontinence O nocturia?
- again--we really need to know what 3rd line therapies are.
- variance of what factors?
- Please include a reference for the interested reader to learn more about these methods .
- so our readers will not get this as explained. Who is voting? (I'm assuming this is the computer model making some sort of assessment but this needs clarification for the O&G readership)
- For clarify, f here in the model development phase is the prediction that a patient will benefit by use of these medications. Correct?
- ROC, AUROC, AUC?
- We prefer to avoid providing p values only unless that is the only appropriate test of significance. Where possible in the abstract AND the text, please provide an effect size (such as an OR or RR) and 95% CI's.
- Sentence 256-260 is quite long. Can you break it up some?
- 3 per night?
- line 270 is somehow incomplete. Can you clarify your meaning?
- please edit sentence 271-275 to be 2 separate sentences.
- Please make sure you clarify in our discussion what the marginal benefit of this is. In your introduction, you note that 50-60% of women get a benefit from these medications so your model predicts accurately about 80% of the time. How will it benefit patients to have this model used if you could predict accurately about 1/2 the time to begin with that they would benefit? Please note that I'm not challenging the benefit, but I'm hoping your discussion will include this information.
- I am a bit confused. would you consider calling your 2 data sets the "Model development data set" and the "external validation data set" throughout your paper? I know it seems clumsy and adds words, but it would add clarity.
- please limit p values to 2 digits given small numbers.
- differ?
- Do you mean of the different AC meds, if they were prescribed anything, they were more likely to have oxybutynin? In the preceding sentence you say they were less likely to have tried a drug, but in the sentence starting on 294 you say they were more likely to receive oxybutynin. Needs to be clearer.
- likelihood of what?
- Perhaps your abstract conclusion (and precis) should be clearer that the model works about 70% of the time overall in predicting failure with best discrimination in women < 40
- Is this actually the bottom line? Correct only 62% of the time? This all needs to be quite clear. How well does it work?
- incorporation OF relatively few....
- risk OF dementia. Please give some statistics to tell us how much the dementia risk is increased.

- 62% prediction of response does not seem reliable. This needs to be tempered.
- you said "fair to good" earlier and that seems more correct and you state this again on line 346.
- again, please temper this
- is "this group" women < 40?
- I think this is a more reasonable description "Fair" for women < 40.
- may potentially BE useful
- where possible, please provide CI's

3. The Editors of Obstetrics & Gynecology are seeking to increase transparency around its peer-review process, in line with efforts to do so in international biomedical peer review publishing. If your article is accepted, we will be posting this revision letter as supplemental digital content to the published article online. Additionally, unless you choose to opt out, we will also be including your point-by-point response to the revision letter. If you opt out of including your response, only the revision letter will be posted. Please reply to this letter with one of two responses:

- A. OPT-IN: Yes, please publish my point-by-point response letter.
- B. OPT-OUT: No, please do not publish my point-by-point response letter.

4. As of December 17, 2018, Obstetrics & Gynecology has implemented an "electronic Copyright Transfer Agreement" (eCTA) and will no longer be collecting author agreement forms. When you are ready to revise your manuscript, you will be prompted in Editorial Manager (EM) to click on "Revise Submission." Doing so will launch the resubmission process, and you will be walked through the various questions that comprise the eCTA. Each of your coauthors will receive an email from the system requesting that they review and electronically sign the eCTA.

Please check with your coauthors to confirm that the disclosures listed in their eCTA forms are correctly disclosed on the manuscript's title page.

5. All submissions that are considered for potential publication are run through CrossCheck for originality. The following lines of text match too closely to previously published works.

Please add some variance to lines 175-181 (the primary outcome...in the analysis). We understand that there will be overlap with previous methods sections, but copying and pasting from a previous paper should be avoided.

6. Standard obstetric and gynecology data definitions have been developed through the reVITALize initiative, which was convened by the American College of Obstetricians and Gynecologists and the members of the Women's Health Registry Alliance. Obstetrics & Gynecology has adopted the use of the reVITALize definitions. Please access the obstetric and gynecology data definitions at <https://www.acog.org/About-ACOG/ACOG-Departments/Patient-Safety-and-Quality-Improvement/reVITALize>. If use of the reVITALize definitions is problematic, please discuss this in your point-by-point response to this letter.

7. Because of space limitations, it is important that your revised manuscript adhere to the following length restrictions by manuscript type: Original Research reports should not exceed 22 typed, double-spaced pages (5,500 words). Stated page limits include all numbered pages in a manuscript (i.e., title page, précis, abstract, text, references, tables, boxes, figure legends, and print appendixes) but exclude references.

8. Specific rules govern the use of acknowledgments in the journal. Please note the following guidelines:

- * All financial support of the study must be acknowledged.
- * Any and all manuscript preparation assistance, including but not limited to topic development, data collection, analysis, writing, or editorial assistance, must be disclosed in the acknowledgments. Such acknowledgments must identify the entities that provided and paid for this assistance, whether directly or indirectly.
- * All persons who contributed to the work reported in the manuscript, but not sufficiently to be authors, must be acknowledged. Written permission must be obtained from all individuals named in the acknowledgments, as readers may infer their endorsement of the data and conclusions. Please note that your response in the journal's electronic author form verifies that permission has been obtained from all named persons.
- * If all or part of the paper was presented at the Annual Clinical and Scientific Meeting of the American College of Obstetricians and Gynecologists or at any other organizational meeting, that presentation should be noted (include the exact dates and location of the meeting).

9. Provide a short title of no more than 45 characters, including spaces, for use as a running foot.

10. Provide a précis on the second page, for use in the Table of Contents. The précis is a single sentence of no more than 25 words that states the conclusion(s) of the report (ie, the bottom line). The précis should be similar to the abstract's conclusion. Do not use commercial names, abbreviations, or acronyms in the précis. Please avoid phrases like "This paper

presents" or "This case presents."

11. The most common deficiency in revised manuscripts involves the abstract. Be sure there are no inconsistencies between the Abstract and the manuscript, and that the Abstract has a clear conclusion statement based on the results found in the paper. Make sure that the abstract does not contain information that does not appear in the body text. If you submit a revision, please check the abstract carefully.

In addition, the abstract length should follow journal guidelines. The word limits for different article types are as follows: Original Research articles, 300 words. Please provide a word count.

12. Only standard abbreviations and acronyms are allowed. A selected list is available online at <http://edmgr.ovid.com/ong/accounts/abbreviations.pdf>. Abbreviations and acronyms cannot be used in the title or précis. Abbreviations and acronyms must be spelled out the first time they are used in the abstract and again in the body of the manuscript.

13. The journal does not use the virgule symbol (/) in sentences with words. Please rephrase your text to avoid using "and/or," or similar constructions throughout the text. You may retain this symbol if you are using it to express data or a measurement.

14. In your Abstract, manuscript Results sections, and tables, the preferred citation should be in terms of an effect size, such as odds ratio or relative risk or the mean difference of a variable between two groups, expressed with appropriate confidence intervals. When such syntax is used, the P value has only secondary importance and often can be omitted or noted as footnotes in a Table format. Putting the results in the form of an effect size makes the result of the statistical test more clinically relevant and gives better context than citing P values alone.

If appropriate, please include number needed to treat for benefits (NNTb) or harm (NNTh). When comparing two procedures, please express the outcome of the comparison in U.S. dollar amounts.

Please standardize the presentation of your data throughout the manuscript submission. For P values, do not exceed three decimal places (for example, "P = .001"). For percentages, do not exceed one decimal place (for example, 11.1%).

15. Please review the journal's Table Checklist to make sure that your tables conform to journal style. The Table Checklist is available online here: http://edmgr.ovid.com/ong/accounts/table_checklist.pdf.

16. Figures

Figure 1: Please consider moving these to supplemental digital content, as these will likely not fit on a printed page. If they should stay in print, then they will need to be split into two separate figures.

Figure 2: Please review n values (746–165 does not equal 582). Additionally, where do the two boxes on the bottom of the page fit in (discharged on postop day 0 and day 1)?

Figure 3: Please upload a high-res version of this figure (eps, tiff, jpeg, etc.). Pasting into Word often results in a lower resolution.

Figure 4: Please upload a high-res version of this figure (eps, tiff, jpeg, etc.). Pasting into Word often results in a lower resolution.

17. Authors whose manuscripts have been accepted for publication have the option to pay an article processing charge and publish open access. With this choice, articles are made freely available online immediately upon publication. An information sheet is available at <http://links.lww.com/LWW-ES/A48>. The cost for publishing an article as open access can be found at <http://edmgr.ovid.com/acd/accounts/ifauth.htm>.

Please note that if your article is accepted, you will receive an email from the editorial office asking you to choose a publication route (traditional or open access). Please keep an eye out for that future email and be sure to respond to it promptly.

18. If you choose to revise your manuscript, please submit your revision through Editorial Manager at <http://ong.editorialmanager.com>. Your manuscript should be uploaded in a word processing format such as Microsoft Word. Your revision's cover letter should include the following:

- * A confirmation that you have read the Instructions for Authors (<http://edmgr.ovid.com/ong/accounts/authors.pdf>), and
- * A point-by-point response to each of the received comments in this letter.

If you submit a revision, we will assume that it has been developed in consultation with your co-authors and that each author has given approval to the final form of the revision.

Again, your paper will be maintained in active status for 21 days from the date of this letter. If we have not heard from you by Aug 15, 2019, we will assume you wish to withdraw the manuscript from further consideration.

Sincerely,

Nancy C. Chescheir, MD
Editor-in-Chief

2018 IMPACT FACTOR: 4.965

2018 IMPACT FACTOR RANKING: 7th out of 83 ob/gyn journals

In compliance with data protection regulations, you may request that we remove your personal registration details at any time. (Use the following URL: <https://www.editorialmanager.com/ong/login.asp?a=r>). Please contact the publication office if you have any questions.

Dear Editors,

We are submitting our revised manuscript entitled "*Development and Validation of a Machine Learning Algorithm for Predicting Response to Anticholinergic Medications for OAB Syndrome*" with changes specifically addressing the reviewers' and editors' comments and critiques. As the lead author, I re-affirm that this manuscript is an honest, accurate, and transparent account of the study being reported; that no important aspects of the study have been omitted; and that any discrepancies from the as planned have been explained. We are only submitting the manuscript to *Obstetrics & Gynecology* and will not submit to any other journal until a final decision is rendered.

Sincerely,

David Sheyn, MD

[Redacted]

[Redacted]

[Redacted]

[Redacted]

Response to Reviewer:

We thank the reviewers for their supportive and insightful comments on our manuscript. We have addressed each of the issues as outlined below. We have restated the comments from the reviewers and highlighted our response in bold/italic font. We have also included the modified text where necessary.

REVIEWER COMMENTS:

Reviewer #1:

Summary: This is a very interesting article that addresses an important clinical question. Often times we treat OAB patients with multiple medications before moving on to 3rd line therapy. With this model, we could potentially triage patients faster and help counsel patients on their risk of treatment failure.

Abstract:

*Line 62, delete the "." after practice

This was deleted.

Methods:

* Can more information be given on the TRIPOD statement .

A short background on TRIPOD was added to the methods section

* Were patients excluded if they had tried an OAB med prior to coming to your office?

Yes. Patients may have been treated with an anti-cholinergic by their PCP or referring provider prior to being seen by a urogynecologist. The "entry" to the study was the initial urogynecologist visit, however, as long as patients have not had any prior third-line treatment and met all other inclusion/exclusion criteria, they were included in the database.

* How was treatment response of > 50% documented.. was this the standard at your institution? Was a validated questionnaire used?

Patient response was determined subjectively. Change in symptom severity was assessed by comparing number of daytime voids, urinary urgency, urgency incontinence episodes and nocturia from initial visit to the first post-treatment follow-up. Patients were also asked to subjectively rate their response in terms of percent improvement from baseline.

Results:

* Line 270: change "never" to no

This was corrected.

* What was the distribution for the other medications prescribed?

Oxybutynin was the most common prescribed medication, 43.1%, followed by trospium, 18.4%, tolterodine, 17.3%, solifenacin, 10.9%, darifenacin, 4.2%, and fesoterodine, 4.7%.

* If the majority were treated with oxybutynin or tolterodine, is your model accurate for all the anticholinergic medications?

Most patients were treated with oxybutynin, 43.1%, with similar distributions of most other medications, tolterodine, trospium and solifenacin. (the reviewer may be referring to the pre-trial entry medication i.e. the "failed medication" in the third paragraph of the results section, which was confusing, and the paragraph was corrected to improve readability). With regard to accuracy of prediction for other anti-cholinergics, there is little evidence to suggest that one anti-cholinergic is better than another in terms of efficacy, and the model performed similarly well on external validation with a different distribution of trialed anti-cholinergic. However, we acknowledge that this is a limitation and this is mentioned in the discussion.

Conclusion:

* Line 321: delete "are" before generally

This was corrected.

* Additional limitation is that all data came from single institution with predominately white and African American population.

This was added as a limitation

* I would also add that no validated questionnaires were used. Most studies these days have a composite score for success, both subjective and objective. This has been shown to have higher correlate with success.

We elaborated upon the use of subjective response as a marker of treatment success being a limitation and lack of using validated questionnaires. However, there is very limited data on whether or not questionnaires are predictive of treatment response, indeed the only questionnaire that has been shown to predict response to medication has been the OABSS, and while the exact wording of the questionnaire was not used in our study, we assessed similar parameters to determine response (change in frequency, urgency, UI, and nocturia episodes). This was also added to the discussion.

Reviewer #2:

1. The objective was clearly defined and relevant to literature for future personalized medicine in treatment of OAB

2. Line 161 discussing follow up: were patients who never reached maximum dosing excluded and if so how do you feel this would impact prediction of all patients starting on treatments for OAB :

Only patients who were non-responders at lower doses and did not try a higher (maximum) dose were excluded; while this may be considered “treatment failure” there would be too many confounders to account for in this scenario (change in insurance coverage, patient frustration etc); patients who achieved treatment response at lower doses were still included in the “success” group.

3. Explain why only ER medication prescriptions were only used for the study inclusion and search-again this could increase the number of inclusion in data analysis and more generalizable data.

IR formulations were included, and patients who responded were termed “responders”, however, patients who did not respond to IR formulations, must have been trialed on an XL/ER formulation at the maximum effective dose prior to the being termed a treatment failure. We now realize that the last sentence of the second paragraph in the methods section does not signify this distinction and have clarified this within the manuscript. These strict criteria for in place in order to minimize the number of false non-responders in the database.

4. Tables need titled and numbered. Difficult to read. Not certain of usefulness to readers of having the Random forests noted in the journal

Tables were edited for clarity and titles were added.

5. Overall commend authors on use of excellent data set and relevant inclusion. Other factors to consider would be pad use, use of depends, and if possible a validated questionnaire at baseline such as MESA scores since this is very subjective and the study in terms of definition of success and failures are also subjective this could lead to understanding if high bothersome is significant in predictive model to failure

Please see response to Reviewer #1. While we did not explicitly utilize a questionnaire for assessment of treatment response, the questions used to determine response for akin to the OABSS which thus far has proven to be the only questionnaire related to treatment response. This is most likely due to lack of investigation of other questionnaires. However, we appreciate the Reviewer’s concern and we agree that an assessment of this model’s prediction accuracy should be compared to those of other measures of symptoms severity and are currently conducting a prospective cohort study to evaluate the concordance between the model and questionnaires.

6. Results: would recommend limited some of the repeat data being reported from Table 3

We agree with the reviewer that some of the tables are redundant, we have consolidated the Table 3 and 4 on the initial submission to a single table (Table 3) which lists the characteristics of both the model development set and external validation set

7. In Discussion: limitation of study should include short follow up median 97 days in data. Clinically when counseling patients, longer term follow up in success would be more relevant -This would not help predict the <20% who continue to use AC after one year which would be more clinically relevant.

We agree that it would be clinically very useful to have a model that would accurately predict which patients will have long term adherence (≥ 1 year). However, the goal of this study was to develop a model that can reliably predict treatment success or failure during the standard medication trial period which is three months in order to spare those patients who are likely to fail treatment an additional 12 weeks of waiting for effective therapy. Anecdotally could be as short as 4 weeks in some FPMRS clinics, depending on practice patterns. Thus we do not think that using a median follow-up of 3 months is a limitation as it reflects current standard clinical practice.

8. Conclusion- again accurate prediction in average follow up of 3 months should be stated

This was added to the conclusion

Reviewer #3:

Abstract - Objective: To develop and externally validate a prediction model for anti-cholinergic response in patient with Overactive Bladder (OAB)

Methods - model retrospective dataset of 559 women with OAB treated with anti-cholinergics between 1/2010-12/17

primary outcome was >50% improvement of patient reported symptoms, groups stratified by age <40, 40-60, >60 and failed AC medications

stratified cross validation with machine learning algorithms - Random Forest (RF) was most accurate treatment failures or responders; RF model externally validated - prospective collected data in 82 women

Results - correctly predicts failure in 81% and response in 61%

Conclusions - validated algorithm can predict AC failure

Intro - OAB incidence, treatment is behavioral and pelvic PT

AC tx - average efficacy 50-60% with <20% continuing after 1 year, goal is to determine which patient benefits from prediction model

objective: develop and validate predictive model to determine likelihood of success or failure of AC treatment using machine learning methods

Materials/ Methods - IRB approval, use of multivariable prediction model for treatment of OAB with AC medication from 1/10-12/17

Patient selection - identified by prescriptions in medical records, inclusion and exclusion criteria are listed

data for 41 clinical risk factors collected with primary outcome being subjective failure defined as no or minimal improvement after a minimum of 4 weeks of treatment on maximum dose with documentation of medication adherence

Model development - patient stratified by age and previous failures

8 features for model, ROC threshold defined, patients classified as responders and non responders

External validation - data on same parameters collected 1/17-12/18 and accuracy/ AUC assessed

Results - 559 patients were followed for 97days - 47.6% with treatment response

accuracy of RF model 80.3% - most predictive for patients <40 and least predictive for patients 41-60 with previous failure

External validation - 82 in external validation, sensitivity and specificity of model with treatment failure as outcome 79.2 and 79.3%

most predictive for patients <40 and least predictive for >60 with previous failure

37.6% responded to treatment and those who didn't respond were correctly predicted in 81%

Discussion - AC medications have a modest improvement, machine learning algorithm uses model to discriminate likelihood of treatment failure

good accuracy in women <40 and >60 if they haven't previously failed

strengths - large sample size with external validation

limitations - retrospective, subjective

developed and validated a model - prospective trials with voiding diaries are needed

Comments:

1) Entire manuscript needs to be thoroughly proofread for multiple errors - grammatical, syntax, and spelling- throughout but particularly in the results and discussion part

2) 209-218 - clarify and proofread model technique - please make more comprehensible

The description of model development was re-worded as to make more comprehensible

3) 240-243 - proofread

This section was reviewed and re-worded

4) why is the model prediction different for the cases than the external validation - what does this reflect?

The difference is likely due to the different patient characteristics between those used in model development and those in the external validation data set. These differences are to be expected as the population of patients treated for OAB will likely differ from our initial group; this would be one of the limitations of our model. For instance, it is unclear how this model would perform in a population composed entirely of Asian, upper socioeconomic status women who are older than 65, as this group is not well represented in the model development set. We have added a functionality to our model that additional patients may be added in order to create a diverse model which should improve prediction accuracy. This function is not yet available for wide-use as it requires HIPPA approval from each site that is entering patients and the model would have to be recalibrated to include new patients and then again internally and externally validated prior to public dissemination.

results - most predictive for patients <40 and least predictive for 41-60 with previous failure but external validation most predictive in <40 and least predictive if >60 with previous failure

5) clarify what a machine learning model is and how it factors in the variables, which variables have which impact.

It would be helpful to know which factors put people at risk for failure

We have elaborated upon how a Random Forest model is developed and some on its functionality, however, we feel that a discussion of machine learning beyond this would be out of the scope of this manuscript. Table 2 lists which risk factors were selected for prediction of response for each patient stratum.

6) The idea of a model is interesting, but if only 37% of the people respond, please indicate what added value this provides - it is already clear that most people won't respond regardless and should be counseled as such

The 37% response rate was only in the external validation set (the patient group that was not used to develop the model). It is expected that if this model is utilized in different clinical settings that the response and failure rates would deviate from what was seen in the model development set (47.6%). It is likely that the model will perform differently with a variation in prevalence of the primary outcome (treatment failure). The response rate of 47.6% is what similar to what is reported in the literature across multiple studies, thus it would be expected that the majority of clinical settings would demonstrate a similar rate of failure. The value of this model would then be primarily to the one-half of the population that is likely to fail medication, allowing for the possibility of bypassing what will be an ineffective drug trial and proceeding directly to more effective therapy. Even in the external validation group with a fairly high prevalence of treatment failure, the model was still able to predict treatment success with an accuracy better than chance, though admittedly not much better, thus it is still capable of discriminating successes from failures even in populations with a high prevalence of failure; furthermore the external validation set had relatively low rate of false positives (patients predicted to fail but actually responding to treatment, a population we wish to keep small), thus it does not uniformly predict that everyone will fail. We feel that with further refinement and addition of more patients, this model will perform quite well in the vast majority of clinical settings, with the

possible exclusion of those with very high rates of treatment failure.

7) are these patients also pursuing pelvic PT - if so, this could confuse the results/responses

This is an excellent point, patients who were undergoing concomitant pelvic floor PT were not included in the analysis.

STATISTICAL EDITOR COMMENTS:

The Statistical Editor makes the following points that need to be addressed:

General: In my version of the article, the Tables and Figures are not labelled, so I will cite them by their numbers given in the text, with, with a short description, as needed.

General and lines 191-198: Since this is not a method our readers would be accustomed to, the Authors should explain how a sufficient sample size vs the proportion of adverse outcomes and how to a priori limit the number of features in the model. That is, how was the number of features in this case chosen as 8? Would a more parsimonious model have similar accuracy and performance? How is the trade-off ascertained between adding more features vs incremental gain in accuracy?

A greater discussion of the F-score and feature selection was added and somewhat simplified.

With regard to a prior selection of feature number The eight selected risk factors were felt to not only independently affect the outcome of treatment failure, but to also have significant interactions between each other, such that the effect of each risk factor on the outcome would be increased or decreased based on the presence or absence of the other selected risk factors.

Modeling with more than eight risk factors was not performed due to concerns regarding over-fitting of the model; exclusion of any of the eight selected risk factors within each patient stratum was not performed due to concerns that this would remove important inter-variable interactions which may influence the outcome prediction.

lines 52-53: Need to clearly identify how many women in the training set and the validation set successfully met the definition of the primary outcome, that is, how many treatment failures or

responders there were overall, and also within the various subsets.

This was clarified in Results section, but not in the abstract due to the word limit. Furthermore, in Table 3 the row initially titled “Responded to Treatment” was changed to “Failed Treatment” to reflect consistency related to the primary outcome

lines 66-71: See later comments re: CIs, for both the training and the validation set.

Confidence Intervals were added where indicated.

Table 3 (Training group characteristics): I presume the hemoglobin A1c were only from the 124 women with DM, but should so state, Should also enumerate all missing values. The last entry "Mean time to first follow-up ...) is formatted as median(IQR), so need to clarify that 97 days represents a median, not a mean value.

The tables were amended to reflect that A1c were related to patients with a diagnosis of diabetes; Table 3 was corrected from “Mean” to “Median”

Table 3 and lines 204-207 (list of RF, SVM and tree augmented Naive Bayes vs 5 combinations of age and NOF): This list of accuracies needs to include CIs, based on the sample sizes. One cannot state that one method is superior because the absolute value of the accuracy is highest. Need to compare statistically, since N = 559 did not represent all women with OAB, but was a sample.

CIs were added where appropriate

lines 280-283: Similarly, the AUCs all need CIs, based on the sample sizes for their subsets by age, and "most predictive" must be based on stats comparisons, not simply which one had the highest arithmetic proportion.

CIs were added where appropriate

lines 298-301: Need to include CIs for the sensitivity and specificity. Again, the comparison of AUCs needs CIs and stats testing. I presume this paragraph is referring to the external validation set only, since that was the subheading on line 285. If this refers to the larger data set, or some combination, should insert a different subheading on line 297.

Yes, this section refers only to the external validation set.

CIs were added where appropriate

lines 303-306: This section's arithmetic seems inconsistent with the total number of women in the test

set and internally inconsistent. For example, since the $N = 82$, then if 31 responded, then the response rate would be 37.8%, not 37.6%. Assuming that 31 responded, then 51 did not, and the model would correctly predict 81.2% of those, which is 41.4. If the overall treatment response was correct 61.5% of the time, that would be 50.4 women. Something is not right. Need to cite internally consistent numbers and CIs. If in fact the correct response of $\sim 61\%$ had a lower bound CI of 50%, then the model was statistically no better than a coin toss.

Thank you for identifying this error; we regrettably must have committed a data entry error, resulting in the incorrect numbers being entered for the false negatives and false positives during calculation of the Sensitivity and Specificity and likelihood ratio, as well as calculating the number of true and false negatives and positives. There were 41 true positives (predicted and actual failures) and 10 that were incorrectly predicted to fail (80.3% correct prediction of failure) and there were 24 true negatives (predicted and actual responders) with 7 who were incorrectly predicted to respond (the correct prediction of response was 77.4%). This resulted in slight changes to Sn (now 80.4%) and Sp (now 77.4%) and the likelihood ratio was 3.56. Since the outcome in question is treatment failure, the correct response was achieved 80.4% of the time; and given the likelihood ratio of 3.56, assuming a pre-test probability of 50% (prevalence of drug refractory OAB), the post-test probability is $\sim 78\%$.

Table 5, lines 286-287: Same issue with "Mean time ..." of mean vs medians. Need to enumerate all missing values and clarify that the hb A1c were only from the DM subsets.

This was corrected; Tables were consolidated and these issues were addressed.

The figures showing decision trees need further explanation. No explanation is given for the gini coefficient, and most of our readers will not be familiar with it, for example. Might be worthwhile to focus on one section of the tree and explain in more detail what is represented.

Upon further discussion between authors and the reviewers' and editorial critiques we believe it is not beneficial to include an example of a decision tree, particularly as it is one of many and serves no other purpose other than an illustration of a decision tree. If the editors feel that it would be valuable to include this image, we would be happy to do so along with a discussion of the gini coefficient, etc.

The figures of AUCs need figure legends and stats corroboration of any statement of superiority. Most of the AUC do not clearly separate, often cross one another and there is no clear hierarchy, except possibly for the highest vs the lowest.

The figures were revised to show each individual AUC.

EDITOR COMMENTS:

1. When you review your paper, I recommend that you read it aloud at some point. Throughout your first submission, there are a number of missing article and pronouns, and incorrect tense use. I've pointed out some of these, but it will perhaps help if you read the paper.

2. Thank you for your submission to Obstetrics & Gynecology. In addition to the comments from the reviewers above, you are being sent a notated PDF that contains the Editor's specific comments. Please review and consider the comments in this file prior to submitting your revised manuscript. These comments should be included in your point-by-point response cover letter.

The notated PDF is uploaded to this submission's record in Editorial Manager. If you cannot locate the file, contact Randi Zung and she will send it by email - rzung@greenjournal.org.

- You will need to include a précis: The précis is a single sentence of no more than 25 words, written in the present tense and stating the conclusion(s) of the report (ie, the bottom line). The précis should be similar to the abstracts conclusion. Do not use commercial names, abbreviations, or acronyms in the précis. Précis should be the "hook" for people who scan the Table of Contents to see what to read. It shouldn't not include statements like "in this study" or "we found". Just state what you found. It may be that the conclusion of your abstract as edited will be good.

A précis was added to the manuscript

- We no longer require that authors adhere to the Green Journal format with the first submission of their papers. However, any revisions must do so. I strongly encourage you to read the instructions for authors (the general bits as well as those specific to the feature-type you are submitting). The instructions provide guidance regarding formatting, word and reference limits, authorship issues, and other things. Adherence to these requirements with your revision will avoid delays during the revision process, as well as avoid re-revisions on your part in order to comply with the formatting.

The author submission guidelines were reviewed, and the manuscript was altered to adhere to the guidelines.

- The way this is worded, I'm unclear if your paper is a secondary analysis of another study or not. I think what you mean is "For the model development, the primary outcome was a >50%...." . Is that correct?

This was clarified in the abstract and body of the manuscript; this is not a secondary analysis

- what is NOF?

Number of Failed (NOF) medications. This acronym was clarified in the abstract and body of the manuscript

- please be consistent. Here you use AUROC; line 66 you use AUC, which is the more common.

AUCROC was changed to AUC

- I'm confused why you used treatment success in the model development phase of your study but treatment

failure in the validation portion. Why not be consistent?

We apologize, this was a writing error; the primary outcome is indeed treatment failure and this was corrected throughout the manuscript.

- likelihood ratio of what?

The likelihood ratio refers to the likelihood of treatment failure, we have corrected this in the manuscript.

- about 80% of the time.

We apologize, but we are not sure what this is referring to.

- Although its not required by the Journal, you. may wish to reference and use the publication guideline for machine learning predictive models. They describe K folding quite well and this may be a useful reference in that regard.

J Med Internet Res. 2016 Dec; 18(12): e323.

Published online 2016 Dec 16. doi: 10.2196/jmir.5870

PMCID: PMC5238707

PMID: 27986644

Guidelines for Developing and Reporting Machine Learning Predictive Models in Biomedical Research: A Multidisciplinary View.

Thank you for this reference, we have added it to the manuscript

- For parsimonious word choice would you consider: "In patients who fail to respond to these treatments, pharmacotherapy is the next option with anti-cholinergic medications the most common choice. "

The sentence was changed to improve readability

- what is 3rd line therapy if not medications? I'm a bit confused by lines lines 123-125. For the 40-50% of women who do not respond to AC treatment, are there other pharmacotherapeutic options? Is it

common to
cycle through the several different AC meds before calling the drug class a failure?

The only other pharmacologic option is myrbetriq which is a sympathomimetic drug, but it has not been shown to be any better than AC therapy; it is rather common for patients to try multiple medications prior to proceeding to third line therapy, often leading to attrition of patients from the practice due to frustration with lack of relief.

- validate A predictive model

This was corrected

- name the IRB here

The name of the IRB was added

- Not sure what you mean by this phrase.

This was clarified

- This is the EMR?

Yes, EMR data was used, this was clarified.

- What if patients were tried on several of these drugs?

Prior to treatment by FPMRS, patients may have been treated by more than one medication by another non-FPMRS provider, this information was collected as "number of previously failed medications, NOF"; Patients would only be prescribed a single drug at a time by an FPMRS physician, and only the initial prescription was considered as the entry point into the study. For instance a patient may have been referred to FPMRS by an internist who had tried oxybutynin and patient did not respond, the patient would then be tried on tolterodine by FPMRS (at this point they are NOF=1 and on tolterodine), the patient fails to respond to tolterodine and is then tried on trospium; at this point the patient is considered a treatment failure (assuming all study criteria are met) and subsequent response to trospium is not assessed.

- For clarity here is it with or without frequency AND with or without urgency AND with or without Nocturia?

OR is it urgency with or without frequency, OR incontinence 0 nocturia?

It is the latter and this was corrected

- again--we really need to know what 3rd line therapies are.

This was clarified.

- variance of what factors?

The definition of the F-score was re-worded

- Please include a reference for the interested reader to learn more about these methods.

A reference was added

- so our readers will not get this as explained. Who is voting? (I'm assuming this is the computer model making some sort of assessment but this needs clarification for the O&G readership)

This was clarified

- For clarify, f here in the model development phase is the prediction that a patient will benefit by use of these medications. Correct?

This was clarified.

- ROC, AUROC, AUC?

This was clarified

- We prefer to avoid providing p values only unless that is the only appropriate test of significance. Where possible in the abstract AND the text, please provide an effect size (such as an OR or RR) and 95% CI's.

p-values are only provided for pairwise comparison of the model development and validation groups; CI were provided for sensitivity, specificity, and AUCs.

- Sentence 256-260 is quite long. Can you break it up some?

This sentence was edited

- 3 per night?

Yes

- line 270 is somehow incomplete. Can you clarify your meaning?

This was corrected.

- please edit sentence 271-275 to be 2 separate sentences.

This was corrected

- Please make sure you clarify in our discussion what the marginal benefit of this is. In your introduction, you note that 50-60% of women get a benefit from these medications so your model predicts accurately about 80% of the time. How will it benefit patients to have this model used if you could predict accurately about 1/2 the time to begin with that they would benefit? Please note that I'm not challenging the benefit, but I'm hoping your discussion will include this information.

An Improved discussion on the benefits of this model has been added.

- I am a bit confused. would you consider calling your 2 data sets the "Model development data set" and the "external validation data set" throughout your paper? I know it seems clumsy and adds words, but it would add clarity.

We appreciate this suggestion and have made the appropriate changes

- please limit p values to 2 digits given small numbers.

Corrected

- differ?

We apologize, but we are not sure what this is referring to. We have gone over the manuscript and have attempted to fix all of the typographical and grammatical errors.

- Do you mean of the different AC meds, if they were prescribed anything, they were more likely to have oxybutynin? In the preceding sentence you say they were less likely to have tried a drug, but in the sentence starting on 294 you say they were more likely to receive oxybutynin. Needs to be clearer.

This was clarified

- likelihood of what?

This refers to the likelihood of treatment failure

- Perhaps your abstract conclusion (and precis) should be clearer that the model works about 70% of the

time

overall in predicting failure with best discrimination in women < 40

This was clarified in abstract and body of manuscript; the overall model accuracy is >80% for treatment failure, the model performs best (AUC=0.83) in women <40 years and worst (AUC=0.70) in women between 40 and 60 years, who have failed one prior medication

- Is this actually the bottom line? Correct only 62% of the time? This all needs to be quite clear. How well does it work?

We have made edits to clarify the bottom line that the model can accurately predict treatment failure

- incorporation OF relatively few....

This was corrected

- risk OF dementia. Please give some statistics to tell us how much the dementia risk is increased.

This was corrected and another reference added

- 62% prediction of response does not seem reliable. This needs to be tempered.

We apologize for the confusion, we were attempting to discuss the PPV and NPV of the model, but did not do so eloquently, we have amended the results section.

- you said "fair to good" earlier and that seems more correct and you state this again on line 346.

- again, please temper this

- is "this group" women < 40?

- I think this is a more reasonable description "Fair" for women < 40.

- may potentially BE useful

- where possible, please provide CI's

The above points were addressed.

3. The Editors of Obstetrics & Gynecology are seeking to increase transparency around its peer-review process, in line with efforts to do so in international biomedical peer review publishing. If your article is accepted, we will be posting this revision letter as supplemental digital content to the published article online. Additionally, unless you choose to opt out, we will also be including your point-by-point response to the revision letter. If you opt out of including your response, only the revision letter will be posted.

Please reply to this letter with one of two responses:

- A. OPT-IN: Yes, please publish my point-by-point response letter.
- B. OPT-OUT: No, please do not publish my point-by-point response letter.

4. As of December 17, 2018, Obstetrics & Gynecology has implemented an "electronic Copyright Transfer Agreement" (eCTA) and will no longer be collecting author agreement forms. When you are ready to revise your manuscript, you will be prompted in Editorial Manager (EM) to click on "Revise Submission." Doing so will launch the resubmission process, and you will be walked through the various questions that comprise the eCTA. Each of your coauthors will receive an email from the system requesting that they review and electronically sign the eCTA.

Please check with your coauthors to confirm that the disclosures listed in their eCTA forms are correctly disclosed on the manuscript's title page.

5. All submissions that are considered for potential publication are run through CrossCheck for originality. The following lines of text match too closely to previously published works.

Please add some variance to lines 175-181 (the primary outcome...in the analysis). We understand that there will be overlap with previous methods sections, but copying and pasting from a previous paper should be avoided.

6. Standard obstetric and gynecology data definitions have been developed through the reVITALize initiative, which was convened by the American College of Obstetricians and Gynecologists and the members of the Women's Health Registry Alliance. Obstetrics & Gynecology has adopted the use of the reVITALize definitions. Please access the obstetric and gynecology data definitions at <https://www.acog.org/About-ACOG/ACOG-Departments/Patient-Safety-and-Quality-Improvement/reVITALize>. If use of the reVITALize definitions is problematic, please discuss this in your point-by-point response to this letter.

7. Because of space limitations, it is important that your revised manuscript adhere to the following length restrictions by manuscript type: Original Research reports should not exceed 22 typed, double-spaced pages (5,500 words). Stated page limits include all numbered pages in a manuscript (i.e., title page, précis, abstract, text, references, tables, boxes, figure legends, and print appendixes) but exclude references.

8. Specific rules govern the use of acknowledgments in the journal. Please note the following guidelines:

- * All financial support of the study must be acknowledged.
- * Any and all manuscript preparation assistance, including but not limited to topic development, data collection, analysis, writing, or editorial assistance, must be disclosed in the acknowledgments. Such acknowledgments must identify the entities that provided and paid for this assistance, whether directly or indirectly.
- * All persons who contributed to the work reported in the manuscript, but not sufficiently to be authors, must be acknowledged. Written permission must be obtained from all individuals named in the

acknowledgments, as readers may infer their endorsement of the data and conclusions. Please note that your response in the journal's electronic author form verifies that permission has been obtained from all named persons.

* If all or part of the paper was presented at the Annual Clinical and Scientific Meeting of the American College of Obstetricians and Gynecologists or at any other organizational meeting, that presentation should be noted (include the exact dates and location of the meeting).

9. Provide a short title of no more than 45 characters, including spaces, for use as a running foot.

10. Provide a précis on the second page, for use in the Table of Contents. The précis is a single sentence of no more than 25 words that states the conclusion(s) of the report (ie, the bottom line). The précis should be similar to the abstract's conclusion. Do not use commercial names, abbreviations, or acronyms in the précis. Please avoid phrases like "This paper presents" or "This case presents."

11. The most common deficiency in revised manuscripts involves the abstract. Be sure there are no inconsistencies between the Abstract and the manuscript, and that the Abstract has a clear conclusion statement based on the results found in the paper. Make sure that the abstract does not contain information that does not appear in the body text. If you submit a revision, please check the abstract carefully.

In addition, the abstract length should follow journal guidelines. The word limits for different article types are as follows: Original Research articles, 300 words. Please provide a word count.

12. Only standard abbreviations and acronyms are allowed. A selected list is available online at <http://edmgr.ovid.com/ong/accounts/abbreviations.pdf>. Abbreviations and acronyms cannot be used in the title or précis. Abbreviations and acronyms must be spelled out the first time they are used in the abstract and again in the body of the manuscript.

13. The journal does not use the virgule symbol (/) in sentences with words. Please rephrase your text to avoid using "and/or," or similar constructions throughout the text. You may retain this symbol if you are using it to express data or a measurement.

14. In your Abstract, manuscript Results sections, and tables, the preferred citation should be in terms of an effect size, such as odds ratio or relative risk or the mean difference of a variable between two groups, expressed with appropriate confidence intervals. When such syntax is used, the P value has only secondary importance and often can be omitted or noted as footnotes in a Table format. Putting the results in the form of an effect size makes the result of the statistical test more clinically relevant and gives better context than citing P values alone.

If appropriate, please include number needed to treat for benefits (NNTb) or harm (NNT_h). When comparing two procedures, please express the outcome of the comparison in U.S. dollar amounts.

Please standardize the presentation of your data throughout the manuscript submission. For P values, do not exceed three decimal places (for example, "P = .001"). For percentages, do not exceed one

decimal place (for example, 11.1%).

15. Please review the journal's Table Checklist to make sure that your tables conform to journal style. The Table Checklist is available online here: http://edmgr.ovid.com/ong/accounts/table_checklist.pdf.

16. Figures

Figure 1: Please consider moving these to supplemental digital content, as these will likely not fit on a printed page. If they should stay in print, then they will need to be split into two separate figures.

Based on reviewer comments we have decided to not include an example of a decision tree in the manuscript, particularly as the ones currently included are just several of many different trees, and do not serve any other purpose other than demonstrating what a decision tree looks like. Unless the editor feels strongly that having a representative decision tree model would be useful to the readers

Figure 2: Please review n values (746–165 does not equal 582). Additionally, where do the two boxes on the bottom of the page fit in (discharged on postop day 0 and day 1)?

This numerical error was corrected. We apologize for the bottom 2 boxes, they were copied and pasted by mistake into this document

Figure 3: Please upload a high-res version of this figure (eps, tiff, jpeg, etc.). Pasting into Word often results in a lower resolution.

Higher resolution images were uploaded

Figure 4: Please upload a high-res version of this figure (eps, tiff, jpeg, etc.). Pasting into Word often results in a lower resolution.

See above

17. Authors whose manuscripts have been accepted for publication have the option to pay an article processing charge and publish open access. With this choice, articles are made freely available online immediately upon publication. An information sheet is available at <http://links.lww.com/LWW-ES/A48>. The cost for publishing an article as open access can be found at <http://edmgr.ovid.com/acd/accounts/ifaauth.htm>.

Please note that if your article is accepted, you will receive an email from the editorial office asking you to choose a publication route (traditional or open access). Please keep an eye out for that future email and be sure to respond to it promptly.

18. If you choose to revise your manuscript, please submit your revision through Editorial Manager at <http://ong.editorialmanager.com>. Your manuscript should be uploaded in a word processing format such as Microsoft Word. Your revision's cover letter should include the following:

- * A confirmation that you have read the Instructions for Authors

(<http://edmgr.ovid.com/ong/accounts/authors.pdf>), and

* A point-by-point response to each of the received comments in this letter.