

RISK ASSESSMENT: LABOR & DELIVERY ADMISSION

Medium Risk

- ☐ Prior cesarean, uterine surgery, or multiple laparotomies
- ☐ Multiple gestation
- ☐ >4 prior births
- ☐ Prior obstetric hemorrhage
- ☐ Large myomas
- ☐ EFW >4000 g
- ☐ Obesity (BMI >40)
- ☐ Hematocrit <30% & other risk

➔ **Type & SCREEN, review protocol**

High Risk

- ☐ Placenta previa/low lying
- ☐ Suspected accreta/percreta
- ☐ Platelet count <70,000
- ☐ Active bleeding
- ☐ Known coagulopathy
- ☐ 2 or more medium risk factors

➔ **Type & CROSS, review protocol**

*** Establish a culture of huddles for high-risk patients & post-event debriefing ***

Appendix 1. Reprinted with permission from American College of Obstetricians and Gynecologists. District II. Safe Motherhood Initiative. Maternal safety bundle for obstetric hemorrhage. Available at: <https://www.acog.org/-/media/Districts/District-II/Public/SMI/v2/HEMSlideSetNov2015.pdf?dmc=1&ts=20190918T1706132151>. Retrieved September 18, 2019. The authors provided this information as a supplement to their article.

RISK ASSESSMENT: INTRAPARTUM

Medium Risk

- [] Chorioamnionitis
- [] Prolonged oxytocin >24 hours
- [] Prolonged 2nd stage
- [] Magnesium sulfate

➔ **Type & SCREEN, review protocol**

High Risk

- [] New active bleeding
- [] 2 or more medium risk factors
(admission &/or intrapartum)

➔ **Type & CROSS, review protocol**

**** Establish a culture of huddles for high-risk patients & post-event debriefing ****

Appendix 1. Reprinted with permission from American College of Obstetricians and Gynecologists. District II. Safe Motherhood Initiative. Maternal safety bundle for obstetric hemorrhage. Available at: <https://www.acog.org/-/media/Districts/District-II/Public/SMI/v2/HEMSlideSetNov2015.pdf?dmc=1&ts=20190918T1706132151>. Retrieved September 18, 2019. The authors provided this information as a supplement to their article.