Appendix 1. Antibiotics Included in Analysis

Penicillins

Amoxicillin
Amoxicillin/clavulanic acid
Ampicillin
Ampicillin/sulbactam
Benxylpenicillin (Penicillin G)
Carbenicillin
Cloxacillin
Dicloxacillin
Methicillin
Nafcillin
Oxacillin
Penicillin
Phenoxymethylpenicillin (Penicillin V)
Piperacillin/tazobactam
Ticaricillin/clavulanic acid

Cephalosporins

First generation

Cephalothin
Cephapirin
Cephradine
Cephalexin
Cefazolin
Cefadroxil

Second generation

Cefamandole
Cefaclor
Cefotetan
Cefoxitin

Duffy CR, Huang Y, Andrikopoulou M, Stern-Ascher CN, Wright JD, Goffman D, et al. Clindamycin, gentamicin, and risk for *Clostridium difficile* infection and acute kidney injury during delivery hospitalizations. *Obstet Gynecol* 2019;134.

The authors provided this information as a supplement to their article.

©2019 American College of Obstetricians and Gynecologists.

Cefuroxime
Cefmetazole
Cefonicid
Cefprozil
Loracarbef

Third generation

Ceftriaxone
Ceftazidime
Cefotaxime
Ceftizoxime
Ceftibuten
Cefixime
Cefperazone
Cefpodoxime
Cefdinir

Fourth generation

Cefepime

Aminoglycosides

Amikacin
Gentamicin
Tobramycin

Carbapenems

Doripenem
Ertapenem
Imipenem/cilastatin
Meropenem

Macrolides

Azithromycin
Clarithromycin

Duffy CR, Huang Y, Andrikopoulou M, Stern-Ascher CN, Wright JD, Goffman D, et al. Clindamycin, gentamicin, and risk for *Clostridium difficile* infection and acute kidney injury during delivery hospitalizations. Obstet Gynecol 2019;134.

The authors provided this information as a supplement to their article.

©2019 American College of Obstetricians and Gynecologists.

Clindamycin
Erythromycin
Fidaxomicin
Spiramycin

Fluoroquinolones

Ciprofloxacin
Levofloxacin
Moxifloxacin
Ofloxacin

Tetracyclines

Doxycycline
Tetracycline

Others

Metronidazole
Vancomycin
Aztreonam
Linezolid
Daptomycin
Chloramphenicol
Sulfamethoxazole/trimethoprim
Nitrofurantin
Actinomycin
Rifampin

The antibiotics in the table were queried in the Perspective database for this analysis along with common trade names.

Duffy CR, Huang Y, Andrikopoulou M, Stern-Ascher CN, Wright JD, Goffman D, et al. Clindamycin, gentamicin, and risk for *Clostridium difficile* infection and acute kidney injury during delivery hospitalizations. Obstet Gynecol 2019;134.

The authors provided this information as a supplement to their article.

©2019 American College of Obstetricians and Gynecologists.

Appendix 2. Demographics and Adjusted and Unadjusted Models for *Clostridium difficile* Infections

	Excluding erythromycin		Excluding metronidazole, vancomycin, and fidaxomicin	
	RR (99% CI)	aRR (99% CI)	RR (99% CI)	aRR (99% CI)
Antibiotics				
No antibiotics	0.04 (0.02, 0.08)**	0.06 (0.03, 0.12)**	0.29 (0.21, 0.40)**	0.58 (0.41, 0.83)**
Clindamycin (+/- other antibiotics)	2.52 (1.87, 3.39)**	2.32 (1.72, 3.13)**	3.50 (2.62, 4.68)**	2.93 (2.19, 3.92)**
Antibiotics excluding clindamycin	Referent	Referent	Referent	Referent
Year				
2006-2008	Referent	Referent	Referent	Referent
2009-2011	0.93 (0.68, 1.27)	0.89 (0.65, 1.21)	0.85 (0.63, 1.15)	0.83 (0.61, 1.11)
2012-2015	1.09 (0.82, 1.45)	1.02 (0.76, 1.36)	1.02 (0.78, 1.34)	0.98 (0.74, 1.29)
Age in years				
18-24	Referent	Referent	Referent	Referent
25-34	1.15 (0.87, 1.52)	1.09 (0.80, 1.47)	1.15 (0.88, 1.50)	1.07 (0.80, 1.43)
35-54	1.60 (1.13, 2.27)*	1.24 (0.85, 1.80)	1.63 (1.16, 2.27)*	1.23 (0.85, 1.77)
Marital Status				
Married	Referent	Referent	Referent	Referent
Single/ Other/Unknown	1.17 (0.92, 1.48)	1.22 (0.93, 1.60)	1.17 (0.93, 1.48)	1.23 (0.95, 1.60)
Race				
White	Referent	Referent	Referent	Referent
Black	1.44 (1.05, 1.97)*	1.19 (0.85, 1.65)	1.42 (1.05, 1.93)*	1.20 (0.87, 1.66)
Other	0.88 (0.67, 1.16)	0.95 (0.71, 1.26)	0.92 (0.70, 1.20)	0.96 (0.73, 1.25)
Payer				
Medicare/Medicaid/ Uninsured/unknown	0.90 (0.71, 1.15)	0.87 (0.67, 1.14)	0.91 (0.73, 1.15)	0.91 (0.70, 1.17)
Commercial	Referent	Referent	Referent	Referent
Preeclampsia				
None	Referent	Referent	Referent	Referent

Duffy CR, Huang Y, Andrikopoulou M, Stern-Ascher CN, Wright JD, Goffman D, et al. Clindamycin, gentamicin, and risk for *Clostridium difficile* infection and acute kidney injury during delivery hospitalizations. *Obstet Gynecol* 2019;134.

The authors provided this information as a supplement to their article.

©2019 American College of Obstetricians and Gynecologists.

Any	4.22 (3.05, 5.85)**	2.11 (1.50, 2.96)**	3.87 (2.80, 5.36)**	2.01 (1.44, 2.81)**
Obstetric comorbidity				
0	Referent	Referent	Referent	Referent
1 or greater	4.30 (3.26, 5.66)**	2.47 (1.85, 3.29)**	4.27 (3.28, 5.56)**	2.55 (1.93, 3.36)**
Mode of delivery				
Vaginal	Referent	Referent	Referent	Referent
Cesarean	5.77 (4.39, 7.58)**	1.90 (1.43, 2.54)**	5.56 (4.29, 7.22)**	3.40 (2.54, 4.56)**
Hospital Teaching				
No	Referent	Referent	Referent	Referent
Yes	1.54 (1.21, 1.95)**	1.39 (1.06, 1.82)*	1.54 (1.23, 1.94)**	1.26 (0.97, 1.63)
Hospital Bed Size				
<400	Referent	Referent	Referent	Referent
400-600	1.24 (0.93, 1.66)	1.08 (0.79, 1.47)	1.35 (1.03, 1.78)*	1.18 (0.88, 1.59)
>600	1.73 (1.30, 2.31)**	1.37 (0.99, 1.88)	1.83 (1.38, 2.41)**	1.42 (1.04, 1.94)*

RR, risk ratio. aRR, adjusted risk ratio. CI, confidence interval. *P<0.01, **P<0.001. The adjusted model included all of the factors in the table.

Duffy CR, Huang Y, Andrikopoulou M, Stern-Ascher CN, Wright JD, Goffman D, et al. Clindamycin, gentamicin, and risk for *Clostridium difficile* infection and acute kidney injury during delivery hospitalizations. Obstet Gynecol 2019;134.

The authors provided this information as a supplement to their article.

©2019 American College of Obstetricians and Gynecologists.

Page 5 of 8

Appendix 3. Demographics and Adjusted and Unadjusted Models For Acute Kidney Injury

Excluding erythromycin		
	Risk ratio (95% CI)	Adjusted risk ratio (95% CI)
Antibiotics		
No antibiotics	0.15 (0.13, 0.18)**	0.34 (0.29, 0.39)**
Gentamicin (+/- other antibiotics)	2.61 (2.26, 3.01)**	2.34 (2.03, 2.70)**
Antibiotics excluding gentamicin	Referent	Referent
Year		
2006	Referent	Referent
2007	1.21 (0.94, 1.56)	1.19 (0.92, 1.53)
2008	1.30 (1.01, 1.67)*	1.23 (0.95, 1.58)
2009	1.62 (1.27, 2.06)**	1.43 (1.12, 1.82)*
2010	1.66 (1.31, 2.11)**	1.20 (0.94, 1.53)
2011	1.59 (1.26, 2.01)**	1.37 (1.09, 1.73)*
2012	1.77 (1.41, 2.22)**	1.46 (1.16, 1.83)**
2013	2.04 (1.64, 2.55)**	1.65 (1.32, 2.06)**
2014	2.13 (1.70, 2.66)**	1.67 (1.33, 2.09)**
2015 1 st quarter	2.27 (1.65, 3.12)**	1.67 (1.21, 2.30)**
Age in years		
18-24	Referent	Referent
25-34	1.03 (0.92, 1.15)	1.08 (0.96, 1.22)
35-39	1.56 (1.35, 1.80)**	1.33 (1.14, 1.55)**
≥ 40	2.81 (2.31, 3.42)**	1.73 (1.41, 2.12)**
Marital Status		
Married	Referent	Referent
Single	1.54 (1.39, 1.70)**	1.26 (1.12, 1.42)**
Other/Unknown	1.65 (1.44, 1.89)**	1.50 (1.29, 1.73)**

Duffy CR, Huang Y, Andrikopoulou M, Stern-Ascher CN, Wright JD, Goffman D, et al. Clindamycin, gentamicin, and risk for *Clostridium difficile* infection and acute kidney injury during delivery hospitalizations. *Obstet Gynecol* 2019;134.

The authors provided this information as a supplement to their article.

©2019 American College of Obstetricians and Gynecologists.

Race		
White	Referent	Referent
Black	2.50 (2.24, 2.81)**	1.62 (1.43, 1.83)**
Other	1.23 (1.10, 1.37)**	1.21 (1.08, 1.36)**
Payer		Referent
Medicare/ Medicaid	1.19 (1.08, 1.30)**	1.09 (0.98, 1.22)
Commercial	Referent	Referent
Uninsured/unknown	1.02 (0.83, 1.25)	1.13 (0.92, 1.40)
Preeclampsia		
None	Referent	Referent
With severe features	44.74 (40.27, 49.70)**	21.57 (19.30, 24.09)**
Without severe features	8.22 (6.97, 9.71)**	5.64 (4.77, 6.66)**
Superimposed	40.32 (34.61, 46.97)**	9.49 (7.97, 11.31)**
Obstetric comorbidity		
0	Referent	Referent
1	5.72 (5.02, 6.52)**	2.22 (1.92, 2.56)**
2	8.53 (7.39, 9.84)**	2.99 (2.58, 3.48)**
>2	21.48 (18.43, 25.04)**	6.79 (5.76, 8.01)**
Mode of delivery		
Vaginal	Referent	Referent
Cesarean	6.28 (5.64, 6.99)**	2.07 (1.83, 2.34)**
Rurality		Referent
Urban	1.90 (1.55, 2.33)**	1.20 (0.97, 1.48)**
Rural	Referent	Referent
Hospital Teaching		
No	Referent	Referent
Yes	2.08 (1.89, 2.28)**	1.33 (1.18, 1.49)**
Hospital Bed Size		

Duffy CR, Huang Y, Andrikopoulou M, Stern-Ascher CN, Wright JD, Goffman D, et al. Clindamycin, gentamicin, and risk for *Clostridium difficile* infection and acute kidney injury during delivery hospitalizations. *Obstet Gynecol* 2019;134.

The authors provided this information as a supplement to their article.

©2019 American College of Obstetricians and Gynecologists.

<400	Referent	Referent
400-600	1.70 (1.53, 1.90)**	1.22 (1.09, 1.37)**
>600	1.97 (1.76, 2.21)**	1.25 (1.09, 1.42)**
Region of Hospital		
Northeastern	Referent	Referent
Midwest	0.90 (0.78, 1.04)	1.01 (0.87, 1.18)
South	0.78 (0.69, 0.88)**	0.78 (0.69, 0.89)**
West	0.53 (0.45, 0.62)**	0.77 (0.65, 0.91)**

CI, confidence interval. *P<0.01, **P<0.001. The adjusted model included all of the factors in the table.

Duffy CR, Huang Y, Andrikopoulou M, Stern-Ascher CN, Wright JD, Goffman D, et al. Clindamycin, gentamicin, and risk for *Clostridium difficile* infection and acute kidney injury during delivery hospitalizations. *Obstet Gynecol* 2019;134.

The authors provided this information as a supplement to their article.

©2019 American College of Obstetricians and Gynecologists.

Page 8 of 8