

**Appendix 1. Characteristics of Women Diagnosed With Endometrial Cancer in March 4-May 26, 2020,
Compared With the Same Period in 2019**

		2019		2020		P value
		N=191	%	N=127	%	
Age, years	18-39	6	3%	7	6%	0.17
	40-49	16	8%	5	4%	
	50-59	45	24%	23	18%	
	60+	124	65%	92	72%	
	Mean	62.4		63		
	Median	63		65		
Race						
	White	99	52%	71	56%	0.82
	Hispanic	25	13%	17	13%	
	African American	18	9%	12	9%	
	Asian or Pacific Islander	33	17%	18	14%	
	Other	15	8%	7	6%	
	Unknown	1	1%	2	2%	
Body mass index (kg/m ²)						
	<18.5	1	1%	1	1%	0.40
	18.5 - <24.9	25	13%	16	13%	
	25- < 29.9	42	22%	23	18%	
	30+	116	61%	76	60%	
	Missing	7	4%	11	9%	
Charlson comorbidity index						
	0	95	50%	57	45%	0.70
	1	36	19%	26	20%	

Suh-Burgmann EJ, Alavi EJ, Schmittiel J. Association of the coronavirus disease 2019 (COVID-19) pandemic with endometrial cancer detection. Obstet Gynecol 2020;136.

The authors provided this information as a supplement to their article.

©2020 American College of Obstetricians and Gynecologists.

	2+	60	31%	44	35%	
Geographic area in Northern California (main counties)						
	North Valley (Sacramento)	40	21%	34	27%	0.32
	East Bay (Alameda, Contra Costa)	34	18%	23	18%	
	South Bay (Santa Clara)	34	18%	13	10%	
	Napa (Napa, Solano)	33	17%	19	15%	
	Central Valley (Fresno)	20	10%	11	9%	
	San Francisco Bay (San Francisco, Marin)	30	16%	27	21%	
Diagnosis by office biopsy (N, %)		160	84%	113	89%	0.19
Diagnosis by dilation and curettage (N, %)		31	16%	14	11%	0.19
Histology* (N, %)						
	Endometrioid grade 1-2	127	66%	81	64%	0.29
	Endometrioid grade 3	22	12%	11	9%	
	Endometrioid grade NOS	5	3%	6	5%	
	Clear cell	2	1%	1	1%	
	Serous	23	12%	20	16%	
	Carcinosarcoma	3	2%	6	5%	
	Other	9	5%	2	2%	

*as determined by endometrial sampling