

Appendix 1. Patient Satisfaction for Nulliparous Women Undergoing Outpatient vs Inpatient Cervical Ripening With a Transcervical Foley Catheter

Outcome	Outpatient n=63	Inpatient n=63	p-value
<u>Survey 1: Six Simple Questions</u> *			
I had appropriate and adequate control over my care	7 (6-7)	7 (7-7)	0.82
The persons responsible for my care are/were caring and compassionate.	7 (7-7)	7 (7-7)	0.72
Problems that have arisen up to now have not been dealt with effectively.	1 (1-2)	1 (1-2)	0.26
My needs have been addressed with appropriate consideration for my time.	7 (7-7)	7 (7-7)	0.99
The overall organization of my care has not been appropriate.	1 (1-1)	1 (1-2)	0.06
I would choose the same type of care for my next pregnancy.	7 (7-7)	7 (6-7)	0.23
<u>Survey 2: Lady-X</u> †			
Presence of expert healthcare professionals during childbirth	1 (1-1)	1 (1-1)	0.30
Information given by the healthcare professionals during childbirth	1 (1-1)	1 (1-1)	0.52

Ausbeck EB, Jauk VC, Files P, Kuper SG, Subramaniam A, Casey BA, et al. Outpatient foley catheter for induction of labor in nulliparous women: a randomized controlled trial. Obstet Gynecol 2020;136.

The authors provided this information as a supplement to their article.

©2020 American College of Obstetricians and Gynecologists.

Taking your wishes seriously during childbirth	1 (1-1)	1 (1-1)	0.47
Emotional support by healthcare professionals during childbirth	1 (1-1)	1 (1-1)	0.19
Feeling of security during childbirth	1 (1-1)	1 (1-1)	0.05
Worries about the health of your child during childbirth	2 (1-2)	1 (1-2)	0.90
Time until the first contact with your child	1 (1-1)	1 (1-1)	0.25
<u>Survey 3: Visual Analog Pain Scale[†]</u>			
Worst amount of pain that you experienced during labor	65 (40-81)	55 (40-80)	0.62
Overall pain that you experienced during labor	60 (40-80)	60 (35-80)	0.79
Worst amount of pain you experienced during placement of the Foley balloon	30 (10-50)	20 (5-40)	0.11
How likely you are to recommend your method of induction to a friend or family member?	80 (50-100)	90 (50-100)	0.45

Data are median (interquartile range) unless otherwise specified.

*Survey scored from 1 to 7 for each question, with 1 indicating “Strongly Disagree” and 7 indicating “Strongly Agree”

†Survey scored from 1 to 3 for each question, with 1 being the most positive response

‡Scored from 0-100 for each question, with 100 being the most positive response

Ausbeck EB, Jauk VC, Files P, Kuper SG, Subramaniam A, Casey BA, et al. Outpatient foley catheter for induction of labor in nulliparous women: a randomized controlled trial. *Obstet Gynecol* 2020;136.

The authors provided this information as a supplement to their article.

©2020 American College of Obstetricians and Gynecologists.