SDC Table 1. OUTLET Study Data Collection	Study Visit							
	Base-	3	6	12	18			
	line	month	month	month	month			
Patient Characteristics								
Demographics (gender, age, race,	X							
education)								
Patient and family income	X							
Pre-injury usual major activities	X							
Motivation to work	X							
Smoking history	X							
Prior leg injury & chronic pain	X							
Pre-injury health related quality of life								
(VR-12)								
Pre-injury activity level (PPAQ)	X							
Marital Status and Social Support	X							
Insurance Information	X							
Co-morbidities, height & weight	X							
Injury Characteristics								
GCS at time of admission	X							
Mechanism	X							
Other non-study injuries	X							
OTA Classification	X							
Gustilo Type or Tscherne Classification	X							
Open Fracture Classification	X							
Extent of muscle, nerve and artery damage	X							
Sensation at time of final closure	X							
Major nerve abnormalities	X							
Foot soft tissue loss	X							
Ipsilateral injuries	X							
Injury x rays (AP and Lateral)	X							
Photos of wound	X				X			
Treatment								
Transfer information (i.e. timing and	X							
mode)								
ICU Admission	X							
First antibiotic use	X							
Discharge disposition	X							
Vascular status	X							
Compartment pressure	X							
Hypotension at time of first surgery	X							
Surgeon survey of factors influencing	X							
treatment decisions								
For Limb Salvage								
Detail on debridement, final fixation, soft	X	X	X	X	X			

	Study Visit							
	Base-	3	6	12	18			
tissue severage revisions and home	line	month	month	month	month			
tissue coverage, revisions and bone								
grafting For Amoutous								
For Amputees	X	X	X	X	X			
Amputation level, type of procedure, wound closure and soft tissue coverage,	Λ	Λ	Λ	Λ	Λ			
revisions and complications								
Clinical Follow-up		X	X	X	X			
Complications related to the study injury		Λ	Λ	Λ	Λ			
Re-hospitalizations (medical record								
review)		v	V	V	V			
Weight Bearing Status & Ambulation		X	X	X X	$\frac{X}{X}$			
Wound & Fracture Healing		X	X					
Pain using VAS & pain medication use		X	X	X	X			
Senation					37			
Range of Motion					X			
Performance Assessments					X			
Patient Reported Outcomes								
Self-Efficacy		X						
Re-hospitalizations (patient reported)		X	X	X	X			
Rehabilitation received		X	X	X	X			
Return to Work/Usual Activity Status		X	X	X	X			
Post traumatic stress disorder (PCL)		X			X			
Depression (PHQ-9)		X			X			
Brief Pain Inventory (BPI)		X			X			
Health Related Quality of Life (VR-12)		X	X	X	X			
Lower extremity function (SMFA)					X			
Paffenbarger Activity Scale (PPAQ)					X			
Phantom pain		X	X	X	X			
Prosthesis use		X	X	X	X			
Satisfaction with Prosthesis (OPUS)					X			

BL=Baseline; PPAQ=Paffenbarger Performance Activity Questionnaire; GCS= Glasgow Coma Score; OTA=Orthopaedic Trauma Association; ICU=intensive care unit; VAS= visual analogue scale; SMFA= Short Musculoskeletal Function Assessment; VR-12 = Veteran's Rand 12 Item Short Form Survey; BPI= Brief Pain Inventory; PHQ= Patient Health Questionnaire; PCL= Post Traumatic Stress Disorder Check List; PSQ= Patient Satisfaction Questionnaire; OPUS = Orthotics Prosthetics Users Survey.