

Supplemental table. Crude odds ratios and 95% confidence intervals for maternal and newborn polymorphisms for whites and blacks separately using the case-control data

	White (n=663)		Blacks (n=227)	
	Mother OR (95% CI)	Newborn OR (95% CI)	Mother OR (95% CI)	Newborn OR (95% CI)
CYP1A1*2A				
-/-*	1.0	1.0	1.0	1.0
-/+	1.03 (0.65-1.66)	0.81 (0.48-1.36)	0.90 (0.51-1.61)	0.56 (0.32-0.99)
+/+	0.50 (0.04-5.60)	∞	0.70 (0.25-1.77)	2.61 (0.68-9.94)
CYP1A2*2B				
-/-*	1.0	1.0	1.0	1.0
-/+ & +/+	1.33 (0.72-2.42)	1.16 (0.64-2.13)	2.91 (0.30-29.50)	0.92 (0.13-6.65)
CYP1A1*4				
-/-*	1.0	1.0	1.0	1.0
-/+ & +/+	1.05 (0.58-1.90)	1.41 (0.73-2.71)	NE	0.22 (0.02-2.03)
GSTT1				
+/+*	1.0	1.0	1.0	1.0
+/-	0.89 (0.63-1.25)	1.47 (1.02-2.09)	0.65 (0.31-1.38)	0.64 (0.28-1.47)
-/-	0.83 (0.53-1.33)	1.27 (0.80-2.03)	0.54 (0.23-1.29)	0.52 (0.20-1.32)
GSTM1				
+/+*	1.0	1.0	1.0	1.0
-/-	0.91 (0.73-1.14)	0.75 (0.54-1.02)	0.75 (0.49-1.15)	1.03 (0.58-1.83)
XRCC3				
Thr/Thr*	1.0	1.0	1.0	1.0
Thr/Met	0.78 (0.55-1.10)	0.84 (0.59-1.18)	1.35 (0.76-2.40)	1.05 (0.61-1.82)

Met/Met	0.77 (0.49-1.19)	0.77 (0.48-1.25)	1.04 (0.20-5.32)	0.95 (0.26-3.45)
---------	------------------	------------------	------------------	------------------

* Reference category

NE indicates that OR could not be estimated; there were 0 cases and 3 controls in the -/+ & +/+ category.