Supplemental Digital Content 5: Proportion of Patients Showing Mean Changes in Area of Lesion and CNV at Baseline and Month 12 (Full analysis set*) (modified last observation carried forward)

	Baseline		Month 12	
	Mean (SD) lesion	Mean (SD)	Mean (SD) lesion	Mean (SD) CNV
	area, mm²	CNV area, mm ²	area, mm²	area, mm²
Ranibizumab 0.5 mg				
(Group I;	4.4.6.70.47	4.04 (4.30)	4.76 (2.22)	0.53 (0.50)
VA stabilization)	4.16 (9.47)	1.04 (1.38)	1.76 (2.22)	0.53 (0.58)
(n = 182)				
Ranibizumab 0.5 mg				
(Group II;	2.00 (4.22)	0.04/1.01	1 02 (2 70)	0.54(0.72)
disease activity)	3.00 (4.22)	0.84 (1.01)	1.83 (2.79)	0.54 (0.73)
(n = 182)				
vPDT				
(Group III)	2.93 (3.98)	0.97 (1.96)	2.10 (2.71)	0.62 (0.67)
(n = 91)				

^{*}Consisted of all patients to whom study treatment were assigned.

SD, standard deviation; VA, visual acuity; vPDT, verteporfin photodynamic therapy