

Online Supplementary File 2: Disease location and behavior (Montreal classification)* in severely, mildly and non anemic children with Crohn's Disease at disease diagnosis (Cohort 2)

	Anemia status		
	Non anemic	Mildly anemic	Severely anemic
Disease location (Montreal Classification)*			
<i>Ileal</i> (L1); n (%)	2 (100)	0 (0)	0 (0)
<i>Colonic</i> (L2); n (%)	4 (25)	8 (50)	4 (25)
<i>Ileo-colonic</i> (L3); n (%)	4 (18)	8 (36)	4 (46)
<i>Isolated upper disease</i> (L4); n (%)	2 (100)	0 (0)	0 (0)
<i>Upper disease colonic</i> (L2+L4); n (%)	5 (23)	6 (27)	11 (50)
<i>Upper disease ileo-colitis</i> (L3+L4); n (%)	9 (19)	27 (56)	12 (25)
Disease behavior (Montreal Classification)*			
<i>Inflammatory</i> (B1); n (%)	23 (23)	44 (45)	31 (32)
<i>Stricturing</i> (B2); n (%)	2 (20)	5 (50)	3 (30)
<i>Penetrating</i> (B3); n (%)	1 (20)	1 (20)	3 (60)

*Silverberg MS et al. 2005 Can J Gastroenterol;19 Suppl A:5-36