

SDC 1: The Proportion of Subjects With OPA Levels Greater Than or Equal to the LLOQ^a (Postvaccination)

| | | % (95% CI ^b) | | | | | | |
|------------------------|----------------|--------------------------|--------------------|---------------------|-------------------|-------------------|---------------------|-------------------|
| Age Group | n ^c | Serotype | | | | | | |
| PCV7 serotypes | | 4 | 6B | 9V | 14 | 18C | 19F | 23F |
| Younger ^d | 175–181 | 100 (98.0–100) | 100 (97.9–100) | 100 (98.0–100) | 100 (97.9–100) | 100 (97.9–100) | 98.9 (96.0–99.9) | 100 (97.9–100) |
| Older ^e | 182–188 | 100 (98.1–100) | 100 (98.0–100) | 100 (98.0–100) | 100 (98.0–100) | 100 (98.0–100) | 100 (98.0–100) | 100 (98.0–100) |
| 6 additional serotypes | | 1 | 3 | 5 | 6A | 7F | 19A | |
| Younger ^d | 175–180 | 97.8 (94.4–99.4) | 99.4 (96.9–100) | 98.9 (96.0–99.9) | 100 (97.9–100) | 100 (97.9–100) | 100 (98.0–100) | |
| Older ^e | 173–187 | 98.9 (96.2–99.9) | 99.4 (97.0–100) | 94.5 (90.2–97.3) | 100 (98.0–100) | 100 (98.0–100) | 100 (98.0–100) | |

^aLLOQ in titers for each serotype was set as follows: serotype 1, 18; serotype 3, 12; serotype 4, 21; serotype 5, 29; serotype 6A, 37; serotype 6B, 43; serotype 7F, 210; serotype 9V, 345; serotype 14, 35; serotype 18C, 31; serotype 19A, 18; serotype 19F, 48; and serotype 23F, 13.

^bExact 2-sided CIs based on the observed proportion of subjects.

^cNumber of subjects with an antibody titer who meet the comparison level for the given serotype.

^d≥5–<10 years of age.

^e≥10–<18 years of age.