Table 2: Healthcare Worker Knowledge of Ebola Transmission and Epidemiology*

True/False Questions asked of HCWs	Total	Physicians	Nurses	p-value
(Correct answer)†	(n=245)	(n=114)	(n=110)	
	# (%)	# (%)	# (%)	
Ebola kills 95% of the people it infects.	124 (51%)	72 (63%)	42 (38%)	<0.001
(False)				
Ebola can be spread by airborne	147 (60%)	81 (71%)	54 (49%)	0.001
transmission. (False)				
The corpses of Ebola patients are not	188 (77%)	94 (82%)	80 (73%)	0.080
infectious. (False)				
The incubation period for Ebola is 3-5	79 (32%)	47 (41%)	24 (22%)	0.002
days. (False)				
A person who has contracted Ebola is	51 (21%)	32 (28%)	12 (11%)	0.001
not infectious until they show symptoms.				
(True)				
Spread via blood (True)	217 (89%)	105 (92%)	95 (86%)	0.165
Spread via urine (True)	153 (62%)	73 (64%)	67 (61%)	0.629
Spread via saliva (True)	185 (76%)	91 (80%)	77 (70%)	0.090
Spread via sweat (True)	116 (47%)	56 (49%)	50 (45%)	0.583
Spread via tears (True)	114 (47%)	54 (47%)	51 (46%)	0.880
Spread via semen (True)	145 (59%)	66 (58%)	66 (60%)	0.749
Spread via gastric secretions (True)	142 (58%)	65 (57%)	64 (58%)	0.860
There are no proven therapies for Ebola.	113 (46%)	69 (61%)	36 (33%)	<0.001
(True)	, ,	, ,	, ,	
Total knowledge score (mean)	56%	61%	50%	0.001

^{*# (%)} represent # of participants answering the questions correctly

†Persons who scored higher on knowledge of Ebola mortality rate and treatment also scored higher on knowledge of how Ebola is transmitted (p=0.001)