

Outcome	Sensitivity	Specificity	Positive predictive value	Negative predictive value	Positive likelihood ratio	Negative likelihood ratio
	%	%	%	%	%	%
<b>CRP &gt;62 mg/L on day 3 or 4</b>						
Hearing threshold >40 dB (better ear)	66	60	53	72	1.65	0.57
Hearing threshold $\geq$ 80 dB	64	50	14	92	1.28	0.72
Any neurological sequelae	71	56	31	88	1.61	0.52
Severe neurological sequelae	83	53	15	97	1.89	0.32
Death	46	50	6	93	0.92	1.08
<b>CRP &gt;62 mg/L on day 3 or 4, and initial Glasgow Coma score &lt;13</b>						
Hearing threshold >40 dB	54	87	71	70	4.15	0.61
Hearing threshold $\geq$ 80 dB	43	76	18	92	1.79	0.75
Any neurological sequelae	55	83	46	87	2.04	0.54
Severe neurological sequelae	78	79	27	97	3.71	0.28
Death	38	75	9	95	1.52	0.83

SDC TABLE 2. Predictive Accuracy of the CRP Values Exceeding the Median on Day 3 or 4, and of the Combination of These CRP Values Combined with the Glasgow Coma Score <13 on Arrival; Analysis of the Latin American Data