

SDC Table 1. Bacterial isolates from 432 blood cultures collected from 309 infants admitted to the neonatal unit, by timing of infection and organism

Twenty cultures grew a significant pathogen and 3 of these cultures were polymicrobial (2 organisms identified) resulting in a total of 23 isolates

Organism	Pre-NU*		NU		Post-NU		Total, n (%)
	CAI†	HAI†	CAI	HAI	CAI	HAI	
<i>Acinetobacter baumannii</i>	0	0	0	1	0	0	1 (4.3)
<i>Acinetobacter</i> sp.	0	0	0	1	0	0	1 (4.3)
<i>Enterobacter cloacae</i>	0	0	0	3	0	0	3 (13.0)
<i>Enterococcus</i> sp.	1	0	0	0	0	0	1 (4.3)
<i>Escherichia coli</i>	0	1	2	0	0	0	3 (13.0)
<i>Klebsiella pneumoniae</i>	1	0	0	6	0	0	7 (30.4)
<i>Moraxella catarrhalis</i>	1	0	0	0	0	0	1 (4.3)
<i>Serratia liquifaciens</i>	0	1	0	0	0	0	1 (4.3)
<i>Serratia marcescens</i>	0	0	0	1	0	1	2 (8.7)
<i>Staphylococcus aureus</i>	0	0	1	0	0	0	1 (4.3)
<i>Streptococcus pyogenes</i>	1	0	1	0	0	0	2 (8.7)
Total, n (%)	4 (17.4)	2 (8.7)	4 (17.4)	12 (52.2)	0 (0.0)	1 (4.3)	23

*NU: neonatal unit

†CAI: community acquired infection; HAI: hospital-acquired infection