

**Table 5. Febrile convulsions up to Month 7**

<b>Vaccine / dose</b>	<b>Age group</b>	<b>Number of vaccine doses<sup>1</sup></b>	<b>Time of occurrence post-vaccination</b>	<b>Intensity</b>	<b>Serious</b>	<b>Causality Assessment<sup>2</sup></b>
IXIARO, 0.25 mL	≥1 to <3 years	2	2 days	Severe	Yes	Not related <sup>3</sup>
IXIARO, 0.25 mL	≥1 to <3 years	2	8 days	Mild	No	Not related <sup>4</sup>
HAVRIX 720	≥1 to <3 years	1	9 days	Severe	Yes	Not related <sup>5</sup>
IXIARO, 0.25 mL	≥1 to <3 years	1	>2 weeks	Moderate	Yes	Not related
IXIARO, 0.25 mL	≥1 to <3 years	2	4 weeks	Severe	Yes	Not related
Prevnar	≥2 months to <1 year	1	4 weeks	Severe	Yes	Not related
HAVRIX 720	≥1 to <3 years	1	4 weeks	Severe	Yes	Not related
IXIARO, 0.25 mL	≥1 to <3 years	2	>5 weeks	Severe	Yes	Not related
IXIARO, 0.25 mL	≥1 to <3 years	2	>7 weeks	Mild	No	Not related
IXIARO, 0.25 mL	≥1 to <3 years	2	>8 weeks	Severe	Yes	Not related
HAVRIX 720	≥1 to <3 years	1	>16 weeks	Moderate	Yes	Not related
IXIARO, 0.25 mL	≥1 to <3 years	2	>5 months	Mild	No	Not related

<sup>1</sup>Vaccine doses received before event onset.

<sup>2</sup>Investigator's assessment of causal association of AE with study vaccination.

<sup>3</sup>Febrile Seizure on the first day of a severe episode of Urinary Tract Infection; no fever on the day of vaccination and following day

<sup>4</sup>Febrile Seizure on the first day of an Upper Respiratory Tract Infection that started 8 days after immunization; no fever on any of the 7 days following immunization

<sup>5</sup>Febrile Seizure on the 3<sup>rd</sup> day of an episode of Acute Tonsillopharyngitis 7 days after immunization; no fever on any of the 6 days following immunization