

Supplemental Digital Content 2 (Table). Differences in Pneumococcal Carriage Prevalence by Age, Sex, Region and HIV status in the Pre-and Post-PCV10 periods, Mozambique, 2012-2015

Characteristic	Pre-PCV10 Introduction			Post-PCV 10 Introduction		
	Total Tested	<i>S. pneumoniae</i> n (%)	P value	Total Tested	<i>S. pneumoniae</i> n (%)	P value
Age, months						
1.4 –11	157	123 (78.3)	reference	176	156 (88.6)	reference
12–23	187	152 (81.3)	0.50	253	214 (84.6)	0.26
24–59	376	304 (80.9)	0.51	482	396 (82.2)	0.05
Sex						
Male	343	283 (82.5)	reference	466	382 (82.0)	reference
Female	377	296 (78.5)	0.18	445	384 (86.3)	0.09
HIV status						
HIV-uninfected	306	242 (79.1)	reference	400	335 (83.8)	reference
HIV-infected	414	337 (81.4)	0.45	511	431 (84.3)	0.86
Region						
Manhiça (Rural South)	501	407 (81.2)	reference	592	506 (85.5)	reference
Nampula (Urban North)	98	82 (83.7)	0.57	150	136 (90.7)	0.11
Maputo (Urban South)	121	90 (74.4)	0.09	169	124 (73.4)	<.001