

Supplemental Digital Content 2: Total plasma concentration-time profiles for (A) ceftolozane and (B) tazobactam, following administration of a single IV dose of ceftolozane/tazobactam, shown as arithmetic mean (±SD). Group 1: ≥12 to <18 years; group 2: ≥7 to <12 years; group 3: ≥2 to <7 years; group 4: ≥3 months to <2 years; group 5: birth (7 days postnatal, >32 weeks gestation) to <3 months.

(A)



- □ Group 1 (Dose 1.5 g)
- Group 2 (Dose 18/9 mg/kg)
- △ Group 3 (Dose 18/9 mg/kg)
- + Group 3 (Dose 30/15 mg/kg)
- × Group 4 (Dose 18/9 mg/kg)
- ∇ Group 5 (Dose 20/10 mg/kg)
- □ Group 6 (Dose 12/6 mg/kg)
- * Group 6 (Dose 20/10 mg/kg)



- □ Group 1 (Dose 1.5 g)
- Group 2 (Dose 18/9 mg/kg)
- △ Group 3 (Dose 18/9 mg/kg) + Group 3 (Dose 30/15 mg/kg)
- × Group 4 (Dose 18/9 mg/kg)
- ▽ Group 5 (Dose 20/10 mg/kg) ☑ Group 6 (Dose 12/6 mg/kg)
- * Group 6 (Dose 20/10 mg/kg)