

Supplemental Digital Content 3

TABLE: Safety and Laboratory Assessments

Assessment	Treatment visits						Follow up visits	
	Baseline	Day 1	Days 2 and 3	Days 4 and ≤15	EOIV	EOT [†]	TOC	LFU
Complete blood count with differential, chemistry panel, CrCL calculation, ESR, and optional CRP	X	If clinically indicated	If clinically indicated	X [‡]	X [§]	If clinically indicated	X [¶]	If clinically indicated
Urine or serum pregnancy test	X						X	
Urine sample routine analysis	X			If clinically indicated				
Intra-abdominal fluid collection for culture	X			If clinically indicated				
Blood sample for culture				If clinically indicated				
Blood for PK analysis ^{††}			X					

[†]Oral therapy only. [‡]Conducted on Day 7 if patient was still on IV study drug at that time. [§]If EOIV occurred within 48 hours after these assessments were performed on study Day 7, they were not repeated. [¶]Performed at TOC only if patient had an abnormal (high/low flag) result on or after EOIV.

^{||}Obtained if clinically indicated and not already collected per standard of care. ^{††}Blood samples were taken in a manner such that the blinded observer remained blinded.

Safety and laboratory tests were performed at a local laboratory at or near the investigator site.

CrCL, creatinine clearance; CRP, C-reactive protein; EOIV, end-of-intravenous treatment; EOT, end-of-treatment; ESR, erythrocyte sedimentation rate; IV, intravenous; LFU, late follow-up; PK, pharmacokinetic; TOC, test-of-cure