

Supplemental Digital Content 11

TABLE: Summary of Baseline Pathogens Isolated in ≥ 2 Patients in Either Treatment Group (micro-ITT Analysis Set)

Pathogen group, n (% of patients) Pathogen	Ceftazidime-avibactam + Metronidazole (N = 50)	Meropenem (N = 19)	Total (N = 69)
Enterobacteriaceae	42 (84.0)	14 (73.7)	56 (81.2)
<i>Escherichia coli</i>	42 (84.0)	13 (68.4)	55 (79.7)
<i>Klebsiella pneumoniae</i>	2 (4.0)	1 (5.3)	3 (4.3)
Gram-negative pathogens other than Enterobacteriaceae	16 (32.0)	10 (52.6)	26 (37.7)
<i>Pseudomonas aeruginosa</i>	14 (28.0)	9 (47.4)	23 (33.3)
Gram-positive pathogens	26 (52.0)	11 (57.9)	37 (53.6)
<i>Streptococcus anginosus</i> group	23 (46.0)	10 (52.6)	33 (47.8)
<i>Enterococcus avium</i>	4 (8.0)	1 (5.3)	5 (7.2)
<i>Enterococcus faecium</i>	2 (4.0)	0	2 (2.9)
Anaerobes	24 (48.0)	12 (63.2)	36 (52.2)
<i>Bacteroides fragilis</i>	14 (28.0)	7 (36.8)	21 (30.4)
<i>Bacteroides fragilis</i> group	2 (4.0)	2 (10.5)	4 (5.8)
<i>Bacteroides caccae</i>	3 (6.0)	0	3 (4.3)
<i>Bacteroides ovatus</i>	2 (4.0)	0	2 (2.9)
<i>Bacteroides thetaiotaomicron</i>	3 (6.0)	3 (15.8)	6 (8.7)
<i>Bacteroides vulgatus</i>	2 (4.0)	0	2 (2.9)
<i>Clostridium perfringens</i>	0	2 (10.5)	2 (2.9)
<i>Clostridium ramosum</i>	2 (4.0)	0	2 (2.9)
<i>Eggerthella lenta</i>	2 (4.0)	0	2 (2.9)
<i>Parabacteroides distasonis</i>	2 (4.0)	0	2 (2.9)
<i>Parvimonas micra</i>	4 (8.0)	5 (26.3)	9 (13.0)
<i>Prevotella buccae</i>	2 (4.0)	0	2 (2.9)

Pathogens were isolated from intra-abdominal site or blood source; n, number of isolates. Multiple isolates of the same species from the same patient were counted only once per pathogen or pathogen group

n, number of isolates. Patients could have more than 1 isolate.