

Table: Potential enteropathogens detected in 185 stools samples from gastroenteritis and 179 stools samples from asymptomatic participating Belgian children.

		Cases	Controls	<i>p</i>
<b>Total</b>		185	179	
<b>Micro-organism(s) found</b>		<b>89/185 (48.1%)</b>	<b>25/179 (14.0%)</b>	< 0.001
<b>Bacteria</b>				
	<i>C. jejuni</i>	<b>26/184 (14.1%)</b>	<b>3/176 (1.7%)</b>	< 0.001
	<i>C. concisus</i>	6/184 (3.3%)	4/176 (2.3%)	NS
	<i>C. coli</i>	0/184 (0.0%)	1/176 (0.6%)	NS
	<i>Salmonella</i>	<b>7/184 (3.8%)</b>	<b>0/177 (0.0%)</b>	0.011
	<i>Aeromonas</i>	7/184 (3.8%)	4/177 (2.3%)	NS
	<i>EPEC</i>	3/184 (1.6%)	3/177 (1.7%)	NS
	<i>Shigella</i>	1/184 (0.5%)	0/177 (0.0%)	NS
	<i>Yersinia</i>	1/184 (0.5%)	1/177 (0.6%)	NS
<b>Virus</b>				
	<i>Adenovirus</i>	<b>11/182 (6.0%)</b>	<b>0/178 (0.0%)</b>	0.002
	<i>Rotavirus</i>	<b>20/182 (11.0%)</b>	<b>0/178 (0.0%)</b>	< 0.001
	<i>Norovirus</i>	8/182 (4.4%)	3/177 (1.7%)	NS
<b>Parasites</b>				
	<i>Blastocystis</i>	5/181 (2.8%)	2/177 (1.1%)	NS
	<i>Dientamoeba</i>	7/181 (3.9%)	3/177 (1.7%)	NS
	<i>Giardia</i>	3/181 (1.7%)	2/177 (1.1%)	NS
	<i>Cryptosporidium</i>	1/181 (0.6%)	0/177 (0%)	NS