

Table E-1 Clinical hip ratings before the operation and at two, five, ten, and fifteen-year intervals

Follow-up interval (yrs)	No. of hips	Rating (percent of hips)				Hip score (points)
		<70 points	70-79 points	80-89 points	90-100 points	
Preop.	100	100	0	0	0	32
2	99	2	4	19	75	93
5	91	2	7	15	76	92
10	77	19	8	18	55	84
15	64	22	25	21	32	81

Table E-2

Evaluation of pain in the hip before the operation and at two, five, ten, and fifteen-year intervals

Follow-up interval (yrs)	No. of hips	Severity of pain (percent of hips)				
		None	Slight	Mild	Marked	Disabling
Preop.	100	0	0	4	69	27
2	99	62	36	2	0	0
5	91	75	23	1	0	1
10	77	90	6	0	4	0
15	64	52	20	16	12	0

Table E-3 Evaluation of pain in the thigh at two, five, ten, and fifteen-year intervals

Follow-up interval (yrs)	No. of hips	Severity of pain (percent of hips)				
		None	Slight	Mild	Marked	Disabling
2	99	81	19	0	0	0
5	91	85	15	0	0	0
10	77	88	10	2	0	0
15	64	88	9	3	0	0

Table E-4 Evaluation of limp before the operation and at two, five, ten, and fifteen-year intervals

Follow-up interval (yrs)	No. of hips	Severity of limp (percent of hips)			
		None	Slight	Moderate	Severe
Preop.	100	0	2	49	49
2	99	50	31	16	3
5	91	59	30	11	0
10	77	34	44	18	4
15	64	33	41	20	6

Table E-5 Evaluation of use of walking aids at two, five, ten, and fifteen-year intervals

Follow-up interval (yrs)	No. of hips	Walking aids (percent of hips)					
		None	Cane for long walks	Cane most of the time	One Crutch	Two Canes	Two crutches or walker
2	99	75	16	9	0	0	0
5	91	71	21	7	0	0	1
10	77	64	25	5	0	0	6
15	64*	53	20	13	2	2	7

\*2% unable to walk secondary to medical reasons at 15 years