

TABLE E-1

Case	Sex	Birth Weight (kg)	Side	Initial Level	Age (mo)	External Rotation (deg)		Recovery of Muscle Group Function at Diagnosis				
						Passive	Loss	Deltoid (C5)	Elbow Flexor (C6)	Triceps (C7)	Finger Flexor (C8)	Intrinsic (T1)
1	M	3.7	R	C5-C6	5	-20	50	2	4	5	5	5
2	M	4.3	R	C5-T1	5	0	60	0	0	0	4	4
3	M	4.1	L	C5-C6	4	-10	70	3	2	5	5	5
4	M	4.8	R	C5-C7	8	0	50	3	4	5	5	5
5	M	3.9	L	C5-T1	6	-20	NA	0	2	2	4	4
6	M	3.9	L	C5-C6	10	-30	NA	3	3	5	5	5
7	F	4.8	L	C5-C6	7	-20	90	1	3	5	5	5
8	F	4.3	L	C5-T1	7	0	NA	0	2	3	1	0
9	F	3.6	R	C5-C7	5	-10	NA	0	2	3	5	5
10	F	4.1	L	C5-C6	3	0	70	2	3	5	5	5
11	F	4.6	R	C5-C6	5	5	40	2	3	5	5	5

In patients seen for the first time with the shoulder already dislocated and fixed in internal rotation, no accurate determination of the amount of the loss of passive external rotation could be made. These patients are included as NA (not available). The strength of recovered muscles was rated as 0 (absent), 1 (trace), 2 (less than antigravity), 3 (greater than antigravity), 4 (strong), and 5 (normal).