

TABLE E1. Clinical Features of the 306 Hips (in 299 patients)

		N	%*
Gender	Female	202	66%
	Male	104	34%
Side	Left	148	48.4%
	Right	158	51.6%
Approach	Anterior	248	81.0%
	Posterior	18	5.9%
	Transtrochanteric	40	13.1%
Fracture Site	Femoral head/neck fractures	205	67.0%
	Inter-/subtrochanteric fractures	75	24.5%
	Acetabulum	23	7.5%
	Not specified	3	1.0%
Fracture Stage	Actual	250	81.7%
	Impending	53	17.3%
	Not specified	3	1.0%
Implant Type	Bipolar	167	54.6%
	Unipolar	55	18.0%
	Total hip	67	21.9%
	Tumor	14	4.6%
	Customized	3	1.0%
Operative Time Intervals	1969-1975	45	14.7%
	1976-1982	61	19.9%
	1983-1989	92	30.1%
	1990-1996	108	35.3%
Complications	Dislocation	11	3.6%
	Periprosthetic fracture	11	3.6%
	Infection	5	1.6%

*Percentages based on the 306 hips.

TABLE E2. Primary Tumor Types

Primary Tumor Types	N	Survival (95% CI)			
		1 Year	2 Years	P Value ⁺	Hazard Ratio (95% CI)
Breast	147	53.7% (46.3-62.4)	30.3% (23.7-38.8)	--	1.0
Kidney	26	26.9% (14.3-50.7)	7.7% (2.0-29.1)	0.04	1.5 (1.0-2.3)
Lung	27	22.2% (11.0-45.0)	11.1% (3.8-32.3)	0.05	1.7 (1.0-2.9)
Prostate	44	40.9% (28.7-58.4)	20.0% (11.0-36.4)	0.10	1.3 (1.0-1.8)
Other	62	20.0% (12.1-33.2)	11.7% (5.8-23.4)	0.01	1.7 (1.1-2.5)

⁺P value derived with use of the Cox proportional hazards regression model.