

Range of Motion Values for Canine Distal (DIP) and Proximal (PIP) Interphalangeal Joints and Flexor Tendon Displacement*

Parameter	Rehabilitation	5 Days		10 Days		21 Days		42 Days	
		4-Strand	8-Strand	4-Strand	8-Strand	4-Strand	8-Strand	4-Strand	8-Strand
PIP Flexion (degree)	Low Force	10.3 ± 3.4	8.5 ± 3.5	8.2 ± 4.9	7.3 ± 3.3	8.4 ± 2.6	8.4 ± 4.8	7.1 ± 2.7	14.1 ± 10.4
	High Force	8.8 ± 3.8	7.2 ± 3.1	12.7 ± 4.5	9.2 ± 3.6	9.0 ± 4.7	9.0 ± 3.0	9.2 ± 4.3	7.8 ± 2.9
DIP Flexion (degree)	Low Force	22.6 ± 5.5	22.1 ± 5.9	24.6 ± 7.4	22.0 ± 4.6	24.0 ± 5.7	27.5 ± 6.7	26.8 ± 6.5	29.0 ± 13.9
	High Force	26.8 ± 2.8	24.7 ± 4.2	23.5 ± 5.5	23.8 ± 4.1	23.8 ± 6.2	29.4 ± 7.7	26.0 ± 4.2	22.8 ± 6.9
Tendon Displacement (mm)	Low Force	4.1 ± 0.5	4.0 ± 1.2	4.9 ± 0.9	4.0 ± 0.8	5.0 ± 0.8	4.4 ± 1.1	5.0 ± 0.9	4.4 ± 0.8
	High Force	4.9 ± 0.7	4.3 ± 0.8	3.9 ± 0.9	4.2 ± 0.9	3.8 ± 0.8	4.6 ± 1.1	4.3 ± 0.7	3.4 ± 0.8

*The values are given as the average and the standard deviation.