

TABLE E-1 Search Strategy

	MEDLINE and Embase	PubMed
Search strategy	1. Knee 2. Tibia 3. Closing wedge 4. Correction accuracy 5. High tibial osteotomy 6. Knee osteotomy 7. Opening wedge 8. Valgus osteotomy knee 9. Corrective osteotomy knee 10. 1 OR 2 11. 3 OR 4 OR 5 OR 6 OR 7 OR 8 OR 9 12. 10 AND 11 13. Limit 12 to (English language)	((((((knee) OR tibia*)) AND ((((((closing wedge) OR correction accuracy) OR high tibial osteotomy) OR knee osteotomy) OR opening wedge) OR valgus knee osteotomy) OR corrective osteotomy knee))) AND English[lang])
No. of papers retrieved	3,140	3,057

References of Included Studies

1. W-Dahl A, Toksvig-Larsen S, Lindstrand A. Ten-year results of physical activity after high tibial osteotomy in patients with knee osteoarthritis. *Knee Surg Sports Traumatol Arthrosc.* 2015 Jul 14 [Epub ahead of print].
2. Bode G, Ogon P, Pestka J, Zwingmann J, Feucht M, Südkamp N, Niemeyer P. Clinical outcome and return to work following single-stage combined autologous chondrocyte implantation and high tibial osteotomy. *Int Orthop.* 2015 Apr;39(4):689-96. Epub 2014 Oct 10.
3. Bonnin MP, Laurent JR, Zadegan F, Badet R, Pooler Archbold HA, Servien E. Can patients really participate in sport after high tibial osteotomy? *Knee Surg Sports Traumatol Arthrosc.* 2013 Jan;21(1):64-73. Epub 2011 Mar 16.
4. Chen LC, Chan YS, Wang CJ. Opening-wedge osteotomy, allografting with dual buttress plate fixation for severe genu recurvatum caused by partial growth arrest of the proximal tibial physis: a case report. *J Orthop Trauma.* 2004 Jul;18(6):384-7.
5. Collins B, Getgood A, Alomar AZ, Giffin JR, Willits K, Fowler PJ, Birmingham TB, Litchfield RB. A case series of lateral opening wedge high tibial osteotomy for valgus malalignment. *Knee Surg Sports Traumatol Arthrosc.* 2013 Jan;21(1):152-60. Epub 2012 Jun 17.
6. Elson DW, Pawelek JE, Shields DW, Dawson MJ, Ferrier GM. Stretching the indications: high tibial osteotomy used successfully to treat isolated ankle symptoms. *BMJ Case Rep.* 2013;2013:20132005. Epub 2013 Sep 10.
7. Faschingbauer M, Nelitz M, Urlaub S, Reichel H, Dornacher D. Return to work and sporting activities after high tibial osteotomy. *Int Orthop.* 2015 Aug;39(8):1527-34. Epub 2015 Feb 26.
8. Hoell S, Suttmoeller J, Stoll V, Fuchs S, Gosheger G. The high tibial osteotomy, open versus closed wedge, a comparison of methods in 108 patients. *Arch Orthop Trauma Surg* 2005 Nov;125(9):638-43.
9. Korovessis P, Katsoudas G, Salonikides P, Stamatakis M, Baikousis A. Medium- and long-term results of high tibial osteotomy for varus gonarthrosis in an agricultural population. *Orthopedics.* 1999 Aug;22(8):729-36.
10. Minzlaff P, Feucht MJ, Saier T, Cotic M, Plath JE, Imhoff AB, Hinterwimmer S. Can young and active patients participate in sports after osteochondral autologous transfer combined with valgus high tibial osteotomy? *Knee Surg Sports Traumatol Arthrosc.* 2016 May;24(5):1594-600. Epub 2014 Dec 7.
11. Niemeyer P, Koestler W, Kaehny C, Kreuz PC, Brooks CJ, Strohm PC, Helwig P, Südkamp NP. Two-year results of open-wedge high tibial osteotomy with fixation by medial plate fixator for medial compartment arthritis with varus malalignment of the knee. *Arthroscopy.* 2008 Jul;24(7):796-804. Epub 2008 Apr 21.
12. Salzmann GM, Ahrens P, Naal FD, El-Azab H, Spang JT, Imhoff AB, Lorenz S. Sporting activity after high tibial osteotomy for the treatment of medial compartment knee osteoarthritis. *Am J Sports Med.* 2009 Feb;37(2):312-8. Epub 2008 Nov 20.
13. Saragaglia D, Rouchy RC, Krayan A, Refaie R. Return to sports after valgus osteotomy of the knee joint in patients with medial unicompartmental osteoarthritis. *Int Orthop.* 2014 Oct;38(10):2109-14. Epub 2014 Jul 4.
14. Schallberger A, Jacobi M, Wahl P, Maestretti G, Jakob RP. High tibial valgus osteotomy in unicompartmental medial osteoarthritis of the knee: a retrospective follow-up study over 13-21 years. *Knee Surg Sports Traumatol Arthrosc.* 2011 Jan;19(1):122-7. Epub 2010 Sep 7.
15. Schaller TM, Roehr B. Salvage of a failed opening wedge tibial osteotomy using a locking plate. *Orthopedics.* 2007 Feb;30(2):161-2.
16. Schröter S, Mueller J, van Heerwaarden R, Lobenhoffer P, Stöckle U, Albrecht D. Return to work and clinical outcome after open wedge HTO. *Knee Surg Sports Traumatol Arthrosc.* 2013 Jan;21(1):213-9. Epub 2012 Jul 10.
17. Takeuchi R, Aratake M, Bito H, Saito I, Kumagai K, Ishikawa H, Akamatsu Y, Sasaki Y, Saito T. Simultaneous bilateral opening-wedge high tibial osteotomy with early full weight-bearing exercise. *Knee Surg Sports Traumatol Arthrosc.* 2008 Nov;16(11):1030-7. Epub 2008 Sep 2.
18. Warme BA, Aalderink K, Amendola A. Is there a role for high tibial osteotomies in the athlete? *Sports Health.* 2011 Jan;3(1):59-69.
19. Waterman BR, Hoffmann JD, Laughlin MD, Burks R, Pallis MP, Tokish JM, Belmont PJ Jr. Success of high tibial osteotomy in the United States military. *Orthop J Sports Med.* 2015 Mar;3(3):2325967115574670. Epub 2015 Mar 12.