COPYRIGHT © BY THE JOURNAL OF BONE AND JOINT SURGERY, INCORPORATED HEIKKINEN ET AL.

AUGMENTED COMPARED WITH NONAUGMENTED SURGICAL REPAIR AFTER TOTAL ACHILLES RUPTURE RESULTS OF A PROSPECTIVE RANDOMIZED TRIAL WITH THIRTEEN OR MORE YEARS OF FOLLOW-UP http://dx.doi.org/10.2106/JBJS.O.00496

Page 1

TABLE E-1 RAND 36-Item Health Survey Scores, Adjusted for BMI, at Fourteen-Year Follow-up Examination

	Mean (SD)			Mean (SD)		
					Finnish Reference	
Domain	Nonaugmented Repair	Augmented Repair	P Value*	Entire Study Group	Population <sup>21</sup>	P Value†
General health	76 (18)	73 (16)	>0.9	75 (17)	58 (7)	< 0.0001
Physical functioning	92 (10)	86 (20)	0.21	89 (16)	81 (7)	0.001
Mental health	87 (9)	86 (13)	>0.9	86 (11)	74 (3)	< 0.0001
Social functioning	95 (12)	95 (9)	>0.9	95 (10)	80 (5)	< 0.0001
Vitality	75 (17)	77 (16)	0.6	76 (17)	64 (5)	< 0.0001
Bodily pain	79 (21)	69 (25)	0.08	73 (24)	71 (6)	0.62
Role functioning/physical	96 (15)	81 (36)	0.08	88 (28)	69 (11)	< 0.0001
Role functioning/emotional	99 (6)	95 (20)	0.7	96 (15)	74 (6)	< 0.0001

<sup>\*</sup>For the difference between the nonaugmented and augmented groups as determined with the BMI-adjusted Student t test. †For the difference between the entire study group and the Finnish reference population as determined with the Student t test.