Copyright © by The Journal of Bone and Joint Surgery, Incorporated Martin et al. Thirty-Day Morbidity After Single-Level Anterior Cervical Discectomy and Fusion: Identification of Risk Factors and Emphasis on ... http://dx.doi.org/10.2106/JBJS.M.00767 Page 1 of 2

TABLE E-1 Patient Inclusion and Exclusion Criteria

Inclusion

CPT codes 22551 and 22554

Exclusion

CPT codes 0095T, 20680, 20970, 22220, 22226, 22522, 22548, 22552, 22585, 22590, 22595, 22600, 22610, 22612, 22614, 22630, 22632, 22808, 22830, 22840, 22841, 22842, 22843, 22846, 22849, 22850, 22855, 22856, 25001, 63015, 63051, 63075 without 22554, 63081, 63082, 63101, 63300, and 63302

Cases for acute trauma

Cases coded with a wound class of 2 or greater

TABLE E-2 Patient Numbers Available for Ea	ch Data Point
Characteristic	No. of Patients Available
Age	2914 (100%)
Female sex	2905 (100%)
Race or ethnicity	2875 (99%)
Body mass index	2903 (100%)
Recent weight loss	2914 (100%)
Diabetes mellitus	2914 (100%)
Current smoker	2914 (100%)
Alcohol use of more than two drinks per day	1952 (67%)
Chronic obstructive pulmonary disease	2914 (100%)
Coronary artery disease	2914 (100%)
Peripheral vascular disease	1952 (67%)
History of transient ischemic attack	1952 (67%)
Dialysis	2914 (100%)
Corticosteroid use	2914 (100%)
Bleeding disorder	2914 (100%)
Preoperative blood transfusion	2914 (100%)
Open wound or wound infection	2914 (100%)
Chemotherapy	1952 (67%)
Radiation therapy	1928 (66%)
Other recent operation	1966 (67%)
Preoperative sepsis	2893 (99%)
White blood cell count	2619 (90%)
Hematocrit	2646 (91%)
Platelets	2621 (90%)
Creatinine	2388 (82%)
Serum albumin	1060 (36%)
International normalized ratio	1626 (56%)
American Society of Anesthesiologists class	2913 (100%)
Operative blood transfusion	793 (27%)
Length of operation	2914 (100%)
Resident involvement	1778 (61%)
Length of stay	2914 (100%)

Copyright © by The Journal of Bone and Joint Surgery, Incorporated Martin et al. Thirty-Day Morbidity After Single-Level Anterior Cervical Discectomy and Fusion: Identification of Risk Factors and Emphasis on . . . http://dx.doi.org/10.2106/JBJS.M.00767 Page 2 of 2

TABLE E-3 Patient Variables Included in the Multivariate Analysis	
Age	
Body mass index	
Diabetes	
History of chronic obstructive pulmonary disease	
Sepsis within thirty days preoperatively	
Dialysis	
Corticosteroid use	
American Society of Anesthesiologists class	
Operative time	
Inpatient or outpatient status	