

TABLE E-1 Patient Demographic Data

Variable	Group 1: Nonsmokers (N = 158)	Group 2: Smokers (N = 69)	Group 3: Former Smokers (N = 24)	P Value
Age* (yr)	30.58 ± 10.97	32.00 ± 9.60	34.75 ± 5.42	0.149
Sex†				0.145
Male	126 (80)	62 (90)	21 (88)	
Female	32 (20)	7 (10)	3 (13)	
Side†				0.675
Left	70 (44)	33 (48)	9 (38)	
Right	88 (56)	36 (52)	15 (63)	
Duration from injury to operation* (mo)	27.54 ± 40.23	27.86 ± 46.15	29.50 ± 37.14	0.977
Additional diagnosis†				0.501
Yes	103 (65)	50 (73)	15 (63)	
No	55 (35)	19 (28)	9 (38)	
*The values are given as the mean and the standard deviation. †The values are given as the number of patients with the percentage of the group in parentheses.				

TABLE E-2 Results of Post-Hoc Testing of Postoperative Variables Between Subgroups of Smokers*

Variable	Light Smokers vs. Moderate Smokers		Light Smokers vs. Heavy Smokers		Moderate Smokers vs. Heavy Smokers	
	P Value	Adj. P Value	P Value	Adj. P Value	P Value	Adj. P Value
SSD in anterior translation	0.442	>0.999	0.011	0.034	0.095	0.284
Lysholm knee score	0.732	>0.999	0.490	>0.999	0.352	>0.999
IKDC subjective score	0.591	>0.999	0.527	>0.999	0.917	>0.999
IKDC objective grade	0.510	>0.999	0.002	0.005	0.102	0.306
*SSD = side-to-side difference. Adj. p value = adjusted p value determined with use of a Bonferroni correction.						