

Patient Satisfaction with pain relief at 2 years for OA, PTOA and RA Groups




Fig. E-1

Patient satisfaction with pain relief at two years for the osteoarthritis (OA), posttraumatic osteoarthritis (PTOA), and rheumatoid arthritis (RA) groups. The x axis presents the percentage of satisfaction.

Patient Satisfaction with Activities of Daily Living (ADL) at 2 years for OA, PTOA and RA Groups




Fig. E-2

Patient satisfaction with activities of daily living at two years for the osteoarthritis (OA), posttraumatic osteoarthritis (PTOA), and rheumatoid arthritis (RA) groups. The x axis presents the percentage of satisfaction.

TABLE E-1 Patient Sex by Group			
Diagnosis	Male (No.)	Female (No.)	Total (No.)
Osteoarthritis	41*	15	56
Posttraumatic osteoarthritis	14	14	28
Rheumatoid arthritis	10	12	22
Total	65	41	106
*P < 0.05.			

TABLE E-2 Adjusted Mean Foot and Ankle Outcome Scores (FAOS) for Pain, Function, and Stiffness by Group				
FAOS	Osteoarthritis	Posttraumatic Osteoarthritis	Rheumatoid Arthritis	P Value
Preoperative scores	(N = 48)	(N = 21)	(N = 19)	
Pain	32.23	36.49	41.86	0.228
Function	41.81	42.65	38.83	0.852
Stiffness	32.32	38.22	35.81	0.602
One-year scores	(N = 33)	(N = 13)	(N = 16)	
Pain	60.65	78.60	78.26	0.042
Function	61.57	76.08	76.37	0.118
Stiffness	57.16	65.38	75.91	0.068
Two-year scores	(N = 28)	(N = 11)	(N = 13)	
Pain	64.72	74.69	77.82	0.297
Function	63.91	69.80	69.30	0.762
Stiffness	52.12	65.47	68.94	0.115