

TABLE E-1 Preoperative Demographic Data

	Group A (N = 86)	Group B (N = 78)	95% Confidence Interval	P Value
Age* (yr)	52.0 (40.1 to 60.3)	51.2 (40.2 to 59.3)	−0.36 to 1.97	0.174
Duration of follow-up* (yr)	10.2 (4.5 to 19.3)	10.7 (5.2 to 17.4)	−1.33 to 0.39	0.282
Male:female ratio (no. of patients)	44:40	39:35		0.968
BMI* (kg/m ²)	26.2 (16.2 to 43.1)	25.9 (16.4 to 42.6)	−1.22 to 1.87	0.567
Lysholm score† (points)	61.2 ± 9.4	62.9 ± 8.6	−4.54 to 1.18	0.249
IKDC score† (points)	60.5 ± 8.8	62.8 ± 8.7	−5.05 to 0.47	0.105
*The values are given as the mean, with the range in parentheses. †The values are given as the mean and the standard deviation.				

TABLE E-2 Clinical Scores and Tibiofemoral Angles at Time of Latest Follow-up*

	Group A (N = 86)	Group B (N = 78)	95% Confidence Interval	P Value
Lysholm score (points)	88.1 ± 7.8	89.4 ± 6.1	−3.53 to 0.95	0.257
IKDC score (points)	90.0 ± 7.9	91.3 ± 5.4	−3.49 to 0.82	0.221
Tibiofemoral angle (°)	3.6 ± 3.2 (varus)	3.2 ± 3.1 (varus)	−0.58 to 1.37	0.427
*The values are given as the mean and the standard deviation.				