					Acetabular	
			Paprosky		Failures and	Dislocations
Study	Hips	Definition	Type (No.)	Follow-up* <i>(yr)</i>	Revisions	(no. [%])
Dearborn and	24	<u>≥</u> 66 mm	Not	7 (5-10.3)	3 infection	5 (21)
Harris <sup>7</sup>			stated		0 loosening	
Hendricks and	12	<u>≥</u> 66 mm	Not	13.9	0 loosening	5 (21)
Harris <sup>8</sup>			stated			
Wedemeyer et	17	<u>≥</u> 64 mm	2 (12)	6.9 (2.7-12.5)	1 infection	1 (5.8)
al. <sup>10</sup>			3A (5)		1 loosening	
Gustke <sup>12</sup>	166	<u>≥</u> 62 female,	2 (119)	6.1	5 revisions	4 (2.4)
		<u>≥</u> 66 male	3 (36)		1 loosening	
Fan et al.11	47	<u>≥</u> 60 female,	1 (6)	5.4 (4-7)	1 infection	5 (10.6)
		<u>≥</u> 64 male	2 (30)		1 loosening	
			3 (11)		1 dislocation	
Patel et al.9	43	<u>≥</u> 62 female,	2 (37)	10 (6-14)	2 loosening	2 (4.6)
		<u>≥</u> 66 male	3 (6)		1 dislocation	
					1 wear	
					1 groin pain	
Whaley et al. <sup>4</sup>	89	<u>≥</u> 62 female,	1 (6)	7.2 (5-11.3)	4 loosening	11 (12)
		<u>≥</u> 66 male	2 (54)		3 dislocation	
			3 (29)		1 infection	
Present study	108†	<u>≥</u> 62 female,	1 (1)	8.1 (2-22)	4 infection	8 (7.4)‡
		<u>≥</u> 66 male	2 (50)		2 loosening	
			3 (57)		3 dislocation	
					2 wear	

\*Values are given as the mean, with the range in parentheses. †Entire cohort, 129. ‡Entire cohort, 12 (10).