Copyright © 2013 by The Journal of Bone and Joint Surgery, Incorporated Cameron et al. History of Shoulder Instability and... http://dx.doi.org/10.2106/JBJS.L.00252

TABLE E-1 Demographics and Instability Outcome by History of Instability at Baseline and for Subjects Who Consented but Did Not Complete Baseline Screening

J	No History at	History of Instability	Consented but No
	Baseline $(N = 588)$	at Baseline $(N = 126)$	Baseline* $(N = 336)$
Age† (yr)	$18.77 \pm 0.96$	18.82 ±1.05	$18.78 \pm 0.98$
Height† (cm)	$176.94 \pm 8.51$	$177.74 \pm 8.52$	$177.98 \pm 7.85$
Weight† (kg)	$74.30 \pm 12.79$	$78.26 \pm 15.73$	$76.85 \pm 13.77$
Sex (no. of subjects [%])			
Male	519 (88.3%)	111 (88.1%)	280 (83.3%)
Female	69 (11.7%)	15 (11.9%)	56 (16.7%)
Instability during follow-up (no.			
of shoulders [%])			
No	1271 (97.6%)‡	103 (87.3%)§	656 (97.6%)
Yes	31 (2.4%)	15 (12.7%)	16 (2.4%)

<sup>\*</sup>Data are for the 336 subjects who consented but were unable to complete all baseline data collection visits due to scheduling conflicts.

<sup>†</sup>The data are presented as the mean plus the standard deviation.

<sup>‡</sup>Column includes 126 uninjured contralateral shoulders from subjects who had a history of instability at baseline.

<sup>§</sup>Column excludes eight shoulders that had a history of surgical stabilization at baseline.