Copyright © by The Journal of Bone and Joint Surgery, Incorporated Li et al. Functional Outcomes After Total Shoulder Arthroplasty in Obese Patients http://dx.doi.org/10.2106/JBJS.L.01145 Page 1 of 1

Perioperative Operating Room Data	BMI Group		
	Normal	Overweight	Obese
ASA classification*	$2.2 \pm 0.6$	$2.1 \pm 0.4$	$2.2 \pm 0.5$
Total intravenous fluid* (mL)	$1446.2 \pm 433.8$	$1375.3 \pm 478.0$	$1687.0 \pm 481.1$
Surgical time from incision to closure* (min)	$108.3\pm19.5$	$115.5 \pm 37.3$	$119.7 \pm 37.3$
No. of operating room complications	None	One glenoid fracture	None

	BMI Group	
Normal	Overweight	Obese
$0.2\pm0.5$	$0.2\pm0.5$	$0.1\pm0.4$
$2.3\pm0.8$	$2.5\pm1.5$	$2.4\pm0.8$
26	22	24
0	3	1
-	0.2 ± 0.5 2.3 ± 0.8	$0.2 \pm 0.5$ $0.2 \pm 0.5$ $2.3 \pm 0.8$ $2.5 \pm 1.5$

<sup>\*</sup>The values are given as the mean and the standard deviation. †One unit of blood was approximately 250 to 300 mL. †The values are given as the number of patients.