

TABLE E-1 Treatment of the Study Groups

	Methicillin-Resistant <i>Staphylococcus aureus</i>		<i>Pseudomonas aeruginosa</i>		
Subgroup	No. of Subjects	Treatment	No. of Subjects	Treatment	Total No.
Bacteriophage	12	1×10^7 Sb-1 bacteriophages in 0.1 mL suspension, locally injected daily for 3 days	12	1×10^7 vB_PsaP PAT14 bacteriophages in 0.1 mL suspension, locally injected daily for 3 days	24
Antibiotic	12	20 mg/kg/day teicoplanin intraperitoneally for 14 days	12	120 mg/kg/day imipenem + cilastatin and 25 mg/kg/day amikacin intraperitoneally for 14 days	24
Bacteriophage + antibiotic	12	1×10^7 Sb-1 bacteriophages in 0.1 mL suspension, locally injected daily for 3 days; and 20 mg/kg/day teicoplanin intraperitoneally for 14 days	12	1×10^7 vB_PsaP PAT 14 bacteriophages in 0.1 mL suspension, locally injected daily for 3 days; and 120 mg/kg/day imipenem + cilastatin and 25 mg/kg/day amikacin intraperitoneally for 14 days	24
Control	12	No treatment	12	No treatment	24
Total	48		48		96