Copyright © 2012 by The Journal of Bone and Joint Surgery, Incorporated Lapner et al. Comparison of Lesser Tuberosity Osteotomy... http://dx.doi.org/10.2106/JBJS.K.01365

TABLE E-1 Baseline Demographic Data

	Peel $(N = 44)$	Osteotomy $(N = 43)$	Total $(N = 87)$
Age* (yr)	$65.3 \pm 11.2 (34.1 \text{ to } 84.7)$	$70.4 \pm 10.1 \ (45.5 \ \text{to} \ 89.7)$	67.8 ± 10.9 (34.1 to
			89.7)
Operative time* (min)	$130 \pm 32 \ (75 \text{ to } 233)$	$140 \pm 43 \ (85 \text{ to } 300)$	135 ± 38 (75 to
			300)
Sex (no. of patients)			
Female	26 (59.1%)	27 (62.8%)	53 (60.9%)
Male	18 (40.9%)	16 (37.2%)	34 (39.1%)
Shoulder affected (no. of patients)			
Left	24 (54.5%)	21 (48.8%)	45 (51.7%)
Right	20 (45.5%)	22 (51.2%)	42 (48.3%)
Arthroplasty type (no. of patients)			
Hemiarthroplasty	9 (20.5%)	7 (16.3%)	16 (18.4%)
Total shoulder arthroplasty	35 (79.5%)	36 (83.7%)	71 (81.6%)
Disease (no. of patients)			
Osteoarthritis	40	42	
Rheumatoid arthritis	4	1	

^{*}The values are given as the mean and the standard deviation, with the range in parentheses.