



Fig. E1

Case 13. Weight-bearing radiographs made preoperatively (top) and one year postoperatively (bottom). The preoperative radiographs illustrate the calcaneofibular impingement (top, left) and the lateral offset on the hindfoot alignment view (top, middle). Postoperatively, the calcaneofibular space is widened and a normalization of the offset of the calcaneus can be seen.

TABLE E-1 Demographic Data, Clinical Manifestation, and Surgical Corrections in Fourteen Patients with Overcorrected Clubfoot Deformity

General				Additional Procedures at Surgery		
Case	Age (yr)	Gender	BMI*	Anterior Closing-Wedge Osteotomy	Calcaneal Osteotomy	First Cuneiform Osteotomy
1	29.1	Female	22.6	No	No	Yes
2	18.8	Female	25.1	Yes	Yes	No
3	43.0	Male	28.1	Yes	No	No
4	28.4	Male	23.6	No	Yes	No
5	65.3	Male	30.8	No	No	No
6	36.4	Male	24.6	Yes	No	No
7	31.4	Male	23.9	No	Yes	No
8	25.2	Male	18.7	No	Yes	No
9	41.2	Female	21.5	No	No	No
10	38.1	Male	25.5	No	No	Yes
11	27.3	Male	30.6	Yes	No	Yes
12	66.0	Male	26.8	Yes	Yes	Yes
13	40.0	Female	45.2	No	No	No
14	26.3	Male	34.4	No	No	Yes
Summary	36.9 ± 14.0*	4 female, 10 male	27.2 ± 6.6*	5 Yes, 9 No	5 Yes, 9 No	5 Yes, 9 No

*BMI = body mass index. †The data are presented as the mean and the standard deviation.