

TABLE E-1 Locations of Soft-Tissue Sarcomas of the Extremity by Anatomical Site	
Anatomical Site	No. of Patients
Upper extremity (N = 100)	
Arm	41 (41.0%)
Shoulder	22 (22.0%)
Hand	15 (15.0%)
Forearm	10 (10.0%)
Scapula	6 (6.0%)
Elbow	4 (4.0%)
Axilla	2 (2.0%)
Lower extremity (N = 263)	
Thigh	134 (51.0%)
Leg	49 (18.6%)
Groin	27 (10.3%)
Ankle	16 (6.1%)
Buttock	14 (5.3%)
Foot	14 (5.3%)
Knee	9 (3.4%)

TABLE E-2 Soft-Tissue Sarcomas of the Extremity Determined by Means of Histopathological Examination	
Histopathological Diagnosis	No. of Patients (N = 363)
High-grade pleomorphic undifferentiated sarcoma*	109 (30.0%)
Leiomyosarcoma	46 (12.7%)
Synovial sarcoma	42 (11.6%)
Liposarcoma	40 (11.0%)
Myxoid liposarcoma	44 (12.1%)
Epithelioid sarcoma	18 (5.0%)
Fibromyxosarcoma	12 (3.3%)
Fibrosarcoma	10 (2.8%)
Spindle cell sarcoma	8 (2.2%)
Other†	34 (9.3%)
<p>*This was formerly known as malignant fibrous histiocytoma. †This category includes alveolar rhabdomyosarcoma, clear cell sarcoma, hemangiopericytoma, lymphangiosarcoma, mixed liposarcoma, pleomorphic liposarcoma, and primitive neuroectodermal tumor.</p>	