**TABLE E-I Patient Characteristics** 

	Group According to Anteroposterior Offset Ratio (AOR)*			
Parameter	AOR = 0	AOR > 0	AOR < 0	P Value
No. (%) of patients	127 (40.1)	103 (32.5)	87 (27.4)	_
Age† (yr)	59.4 (35.3 to 82.9)	58.5 (27.6 to 79.9)	58.3 (25.3 to 85.3)	0.768‡
Sex (M:F)	65:62	51:52	50:37	0.518§
Side (R:L)	65:62	65:38	36:51	0.011§
Bodymass index† $(kg/m^2)$	26.8 (19.4 to 39.5)	26.8 (19.6 to 39.2)	27.3 (19.1 to 39.4)	0.658‡
Etiology of ankle osteoarthritis (no. [%])				0.323§
Posttraumatic	102 (80.3)	91 (88.4)	72 (82.8)	
Primary	18 (14.2)	9 (8.7)	8 (9.2)	
Inflammatory	7 (5.5)	3 (2.9)	7 (8.0)	
ASA classification# (no. [%])				0.981§
1	94 (74.0)	80 (77.7)	66 (75.9)	
2	30 (23.6)	21 (20.4)	19 (21.8)	
3	3 (2.4)	2 (1.9)	2 (2.3)	
4	_	_	_	
Diabetes mellitus (Y:N)	9:118	6:97	5:82	0.897§

<sup>\*</sup>Group assignments were made with use of the AOR as measured from lateral radiographs at time of latest follow-up. †Data are presented as the mean, with the range in parentheses. ‡Analysis of variance test. \$Chi-square test. #ASA = American Society of Anesthesiologists.