TABLE E-1 Inclusion and Exclusion Criteria

TABLE E-1 Inclusion and Exclusion Criteria				
Inclusion Criteria	Exclusion Criteria			
Published prospective series	Case studies			
describing outcomes of primary				
ACL reconstruction				
Five-year minimum follow-up	Less than five-year			
	minimum follow-up			
Reconstruction with patellar	Utilization of graft tissue			
tendon or hamstring tendon	other than patellar tendon or			
autograft	hamstring tendon			
Utilization of all-arthroscopic,	Open ACL reconstruction			
arthroscopic-assisted, or mini-				
arthrotomy technique				
Provide data on both graft	Multiple ligamentous knee			
failure rates and contralateral	injuries other than low-			
ACL rupture rates	grade MCL injuries*			
	Inclusion of skeletally			
	immature patients			
	Use of allograft			
	Animal studies			
	In vitro studies			
	Non-English-language			
	studies			
	Reviews without original			
	data			
	Inclusion of revision ACL			
	reconstructions			
	Use of artificial ligaments			
	Retrospective studies			
*MCI 1' 1 11 / 11'				

<sup>\*</sup>MCL = medial collateral ligament.

TABLE E-2 Rehabilitation Milestones

		Time to Partial Weight-	Time to Full Weight-	
Author	Year	Bearing	bearing	Brace Use
Deehan et al. <sup>5</sup>	2000	Immediate	Immediate	None
Drogset and	2002	Immediate	Immediate	None
Grøntvedt <sup>6</sup>				
Keays et al. <sup>10</sup>	2007	Immediate	Immediate	Extension brace $\times$ 1-2
				weeks
Roe et al. 16	2005	Immediate	Immediate	None
Sajovic et al. <sup>18</sup>	2006	Immediate	Immediate	Extension brace $\times$ 3
				weeks
Shelbourne and	2009	Immediate	Immediate	None
Gray <sup>19</sup>				