

TABLE E-1 Demographic and Amputation Data on 137 Limbs in 123 Patients

Variable of Interest	Entire Study (N = 137)	Burgess (N = 100)	Bone Bridge (N = 37)	P Value
Mean age (<i>yr</i>)	24.7	25.2	23.3	0.01*
Time until arrival (<i>d</i>)	5	5	5	0.06
Follow-up time (<i>d</i>)	748	811	725	0.20
Late amputations† (%)	20	12	41	0.0002*
Amputation in zone of injury (%)	67	77	41	<0.0001*
Mean residual limb length (<i>cm</i>)	14.3	13.27	16.94	<0.0001*

*Denotes significant difference. †Late amputations are defined as those performed ≥ 45 days after injury.

TABLE E-2 Secondary Outcome Measures

Variable of Interest	Entire Study* (N = 137)	Burgess* (N = 100)	Bone Bridge* (N = 37)	P Value
Able to walk	129 (94%)	95 (95%)	34 (92%)	0.69
No assistance required	106 (77%)	76 (76%)	30 (81%)	0.76
Managed with narcotics	44 (32%)	32 (32%)	12 (32%)	1.00
Managed with neuropathic medication	46 (33%)	33 (33%)	13(35%)	0.81
Returned to duty	21 (15%)	13 (35%)	8 (22%)	0.21